

Trajectories of women's physical and psychosocial health following obstetric fistula repair in Uganda: a longitudinal study

Alison M. El Ayadi¹, Justus Baragine², Abner Korn¹, Othman Kakaire², Janet Turan³, Susan Obore⁴, Josaphat Byamugisha², Felicia Lester¹, Hadija Nalubwama², Haruna Mwanje⁴, Vandana Tripathi⁵ and Suellen Miller¹

¹ Department of Obstetrics, Gynecology and Reproductive Sciences, Bixby Center for Global Reproductive Health, University of California, San Francisco, CA, USA

² Department of Obstetrics and Gynaecology, Makerere University College of Health Sciences, Kampala, Uganda

³ Department of Health Care Organization and Policy, School of Public Health, University of Alabama at Birmingham, Birmingham, AL, USA

⁴ Urogynaecology Division, Mulago National Referral and Teaching Hospital, Kampala, Uganda

⁵ EngenderHealth, New York, NY, USA

Abstract

OBJECTIVES To explore trajectories of physical and psychosocial health, and their interrelationship, among women completing fistula repair in Uganda for 1 year post-surgery.

METHODS We recruited a 60-woman longitudinal cohort at surgical hospitalisation from Mulago Hospital in Kampala Uganda (Dec 2014–June 2015) and followed them for 1 year. We collected survey data on physical and psychosocial health at surgery and at 3, 6, 9 and 12 months via mobile phone. Fistula characteristics were abstracted from medical records. All participants provided written informed consent. We present univariate analysis and linear regression results.

RESULTS Across post-surgical follow-up, most women reported improvements in physical and psychosocial health, largely within the first 6 months. By 12 months, urinary incontinence had declined from 98% to 33% and general weakness from 33% to 17%, while excellent to good general health rose from 0% to 60%. Reintegration, self-esteem and quality of life all increased through 6 months and remained stable thereafter. Reported stigma reduced, yet some negative self-perception remained at 12 months (mean 17.8). Psychosocial health was significantly impacted by the report of physical symptoms; at 12 months, physical symptoms were associated with a 21.9 lower mean reintegration score (95% CI –30.1, –12.4).

CONCLUSIONS Our longitudinal cohort experienced dramatic improvements in physical and psychosocial health after surgery. Continuing fistula-related symptoms and the substantial differences in psychosocial health by physical symptoms support additional intervention to support women's recovery or more targeted psychosocial support and reintegration services to ensure that those coping with physical or psychosocial challenges are appropriately supported.

keywords vesicovaginal fistula, rectovaginal fistula, obstetric labour complications, urinary incontinence, social stigma, quality of life

Introduction

Prolonged obstructed labour combined with delays in accessing comprehensive emergency obstetric care results in multifaceted physical trauma, often involving multiple organ systems, termed obstructed labour injury complex [1]. Although all the conditions associated with obstructed labour injury complex represent significant maternal morbidity, obstetric fistula has been the primary focus of public

health programming due to its severe physical, psychological and social sequelae [2–6]. Women with fistula experience constant leakage of urine and/or faeces, are subjected to stigma that frequently results in familial and community isolation [7, 8], and report high rates of depression [6, 9–12]. Accurately estimating the burden of obstetric fistula is a challenge; estimates suggest that as many as 2 million women globally may be affected, with thousands of annual incident cases [13–15].

Fistula closure rates at hospital discharge range from 65% to 90% [16–18]. Due to the array of physical and psychosocial correlates of fistula, longitudinal studies that follow women beyond the period of short-term clinical outcomes are of significant interest [3, 19, 20]. Studies assessing patient-focused outcomes of surgery have identified significant short-term improvements including enhanced overall quality of life [21], psychological status [22], social functioning [23, 24] and economic status [25]. However, they also report significant long-term challenges, including persistent residual incontinence [20, 26, 27], pain and weakness [21, 28], and sexual and fertility complications [5, 21, 29–31]. Lingering physical problems such as these reduce women's ability to resume social roles, making them less likely to consider themselves to have recovered [3, 21, 25, 32,] and placing them at greater risk of continued poor mental health [28, 33, 34].

Most longitudinal studies of fistula recovery are limited to a single, short-term, post-surgical follow-up, precluding our ability to understand the process of longer-term physical and psychosocial recovery. Given the significant physical and psychosocial morbidities experienced by women with fistula, and the lengthy period of time that many women live with fistula prior to accessing treatment, physical and psychosocial recovery is likely a lengthier and more complex process than can be captured at one time point. This study explores women's trajectories of physical recovery and their relationship to psychosocial health following surgical repair of obstetric fistula in Uganda, assessed at four time points over 12 months post-surgery.

Methods

We recruited a longitudinal cohort of women accessing fistula surgery at Mulago National Referral and Teaching Hospital in Kampala, Uganda from December 2014 to June 2015 [35]. Fistula repair within the urogynaecology division is an ongoing surgical service supplemented by several annual fistula repair camps.

Women with obstetric or childbirth-related iatrogenic fistula were eligible for participation upon completion of initial examination and clearance for surgery if they spoke Luganda or English, resided in a community with cellular telephone coverage, and were capable of providing informed consent. Patients scheduled for surgery were approached and screened for eligibility by research staff, who obtained written informed consent or thumbprint confirmation. Where eligible participants were unable to be approached prior to surgery, they were asked to participate post-operatively, following sufficient recovery.

Non-enrolled eligible women included one woman whose fistula resolved through urinary catheterisation and another who refused surgery. Data were captured from the participants prior to surgery (baseline), and 3, 6, 9, and 12 months post-surgery, and abstracted from medical records. Our baseline questionnaire was administered in person by local research staff in the participant's preferred language and included questions on socio-demographic characteristics, obstetric history, and physical and psychosocial health measures (Table S1). Aetiology of fistula was inconsistently captured in medical records; we classified fistula aetiology into obstetric *vs.* likely childbirth-related iatrogenic using criteria proposed by Raasen *et al.* [36]. Subsequent surveys included questions on changeable socio-demographic characteristics, and physical and psychosocial health measures. All four post-discharge follow-up surveys were administered over mobile telephone; telephones and airtime were provided to participants. Physical health assessment measures included the Stanford Self-Rated Health measure for general health [37], the International Consultation on Incontinence Questionnaire Short Form (ICIQ-SF) [38], self-reported experience of other fistula-related symptoms (i.e., faecal incontinence, general body weakness, general pain, pain with urination, vaginal pain, skin irritation, vaginal discharge and difficulty walking) and menstrual irregularity. Psychosocial health assessment included a post-surgical reintegration success instrument, operationalised as global functioning status specific to women affected by fistula [39], the Hopkins Symptom Checklist (HSC) for depression [40, 41], the WHO Quality of Life (QOL)-BREF for quality of life [42, 43], the Rosenberg self-esteem scale [44], a fistula-related stigma assessment modified from a scale used to measure HIV-related stigma [45], the Primary Care Post-traumatic Stress Disorder (PTSD) Screen for trauma [46] and the multidimensional scale of perceived social support [47, 48]. We considered a mean HSC score of >1.75 as positive for depressive symptomatology [49, 50]. The QOL-BREF and PTSD screen were not administered at 3 or 9 months to reduce participant response burden. We defined depressive symptomatology using mean HSC score of 1.75 or above, based on commonly used criteria [51, 52]. Each interview lasted between 30 and 60 min. Medical record abstraction captured fistula characteristics and surgical outcomes.

Socio-demographic and obstetric characteristics were described using univariate analyses: means and standard deviations (SD) or medians and interquartile ranges (IQR) for continuous variables and proportions across categorical variables. Physical and psychosocial health measures were standardised to range 0–100 for

comparability and are presented similarly. Analyses of psychosocial health separately evaluated domains comprising the WHO QOL-BREF (domains: overall, physical, environment, social relationships and psychological) and the fistula-related stigma assessment (domains: negative self-perception, social isolation, verbal abuse and fear of contagion). We estimated separate linear regression models to understand the relationship between any persistent physical symptoms over time (urinary incontinence, faecal incontinence, general body weakness, pain, pain with urination, vaginal pain, skin irritation, vaginal discharge and difficulty walking) and our psychosocial health measures of reintegration, self-esteem, depression, quality of life and stigma. All analyses were performed with Stata v14 software (StataCorp, College Station, TX, USA). Differences were considered statistically significant at $P < 0.05$.

The study protocol was approved by the Makerere University College of Health Sciences, School of Medicine Research and Ethics Committee, the Uganda National Council for Science and Technology, and the University of California, San Francisco Human Research Protection Program, Committee on Human Research.

Results

Socio-demographic and obstetric characteristics of study participants

Research staff screened a total of 79 individuals; 62 were eligible, and 60 were enrolled. Retention was 97% at 12 months post-surgery. Median age at study entry was 28 years (IQR 21–36) (Table 1). Educational attainment varied, but most women had not completed primary school (67%), nor did they work outside the home (58%). Primary sources of financial support were self (30%), husbands (35%) or other individuals (35%), primarily relatives. Household assets varied, with most participants reporting having a radio (58%) and a mobile phone (65%) in the household. Two-thirds of study participants had living children (65%).

Median age at first birth was 18 years (IQR 17–20). Most women had vesicovaginal fistula only (95%), two women had both vesicovaginal and rectovaginal fistula (3%), and one woman had rectovaginal fistula only (2%). Forty per cent of our participants had developed fistula at their first delivery (40%), with 52% reporting a prior live birth. Seven women (12%) had delivered since fistula development. Time between fistula development and surgery ranged from 2 weeks to 31 years: 13% accessed surgery within 1 month, and 32% between 1 and 3 months; however, 28% lived with fistula for more

Table 1 Socio-demographic and obstetric characteristics of study participants at baseline ($n = 60$)

Characteristic	<i>n</i>	%
Current age in years (median, IQR)	28 (21–36)	
Living situation		
Alone*	13	31.6
Husband*	24	40.0
Parents*	8	13.3
Other*	15	25.1
Any living children	39	65
Educational attainment		
None	10	16.7
Some primary	24	40.0
Completed primary	17	28.3
Any secondary	9	15
Occupation		
None	35	58.3
Vendor/shopkeeper	5	8.3
Farmer	15	25.0
Other	5	8.3
Primary source of financial support		
Self	18	30
Husband	21	35
Other	21	35
Household assets		
Piped water	9	15.0
Flush/pour flush toilet	4	6.7
Electricity	26	43.3
Radio	35	58.3
Television	17	28.3
Mobile phone	39	65
Refrigerator	26	43.3
Age at first birth†	18 (16.5–20)	
Pregnancies before fistula†	1 (0–3)	
Live births before fistula†	1 (0–3)	
Any births since fistula†	7	11.7
Age at fistula†	22.5 (18–31)	
Duration living with fistula		
<1 month	8	13.3
1–3 month	19	31.7
3–12 month	9	13.3
1–2 years	3	5
3–5 years	5	8.3
>5 years	17	28.3
Any ANC for pregnancy resulting in fistula	56	93.3
Health facility delivery for pregnancy resulting in fistula	58	96.7
Infant survived delivery for pregnancy resulting in fistula	17	28.3
Likely aetiology of fistula		
Obstetric	41	68.3
Childbirth-related iatrogenic	19	31.7

Iqr, interquartile range.

*With or without young children.

†Maximum two participants per each 'other' district.

than 5 years. Most women delivered at a health facility (97%), and only 28% of infants survived. Nearly, one-third of fistula were classified as likely childbirth-related iatrogenic (31.7%).

Physical health across study follow-up

Most women reported fistula-related symptoms at baseline (Figure 1; Table 2). Self-report of general health status was poor (82%) or fair (18%). Most women reported urinary incontinence (98%), with 75% reporting it all of the time, and 20% reporting it several times per day. Three women reported faecal incontinence (5%). Nearly half of women reported vaginal pain (45%) and 37% reported general pain. Approximately one-third reported weakness (33%), difficulty walking (33%), vaginal discharge (32%) or skin irritation (28%). At hospital discharge, leaking was resolved among 37 of 59 women with VVF (63%) and two of three women with RVF. Of the 22 leaking urine at hospital discharge, nine had repair breakdown only, 10 had post-repair urethral incontinence only and one had both repair breakdown and post-repair urethral incontinence. The origin of leakage for two participants was not available in their medical record.

Across the 12-month study, most women reported improvements in physical health (Table 2). The largest improvement in general health status was observed from baseline to 3 months where the report of general health as excellent, very good or good increased from 0% to 60%. Continued improvement was observed through 6 months (71%). Urinary incontinence, reported by 98% at baseline, was the most prevalent continuing physical symptom. The greatest improvement in urinary incontinence was between baseline and 3 months (39%). Continued reduction in urinary incontinence was reported through 6 months (33%). Among the 19 women reporting persistent incontinence at 12 months, 53% reported it at least once to several or more times per day, and 26% reported it once per week or less. Among those with faecal incontinence at baseline ($n = 3$), the one post-surgical persistent case resolved by 6 months. Among women aged <45 years, 64% reported amenorrhoea at baseline, which dropped to 43% at 3 months, and to 28% at 6 months.

Other fistula-related physical symptoms also declined over follow-up (Figure 1). Large declines were observed from baseline through 3 months in vaginal pain (45–15%), general pain (37–12%), difficulty walking (33–12%) and skin irritation (28–9%). Continued but less pronounced reductions were observed for vaginal pain, general pain and difficulty walking. By 12 months after

surgery, 9% reported general pain, 7% skin irritation, 5% reported vaginal pain and 4% difficulty walking. Declines in vaginal discharge were less steep than the other symptoms, dropping from 32% to 20% in the first 3 months following surgery, with only 2% reporting vaginal discharge by 12 months. Pain with urination dropped rapidly between baseline and 3 months (30% to 19%), plateaued at 6 through 9 months (15%) and further decreased by 12 months (7%). General weakness changed minimally from baseline to 3 months (33–27%) and declined more steeply through 6 months (9%), but this gain was not maintained; general weakness at 9 and 12 months was 16% and 17%, respectively. After urinary incontinence, weakness was the second most prevalent fistula-related symptom at 12 months post-surgery.

Reintegration and psychosocial health across study follow-up


Similarly, improvement was observed across study follow-up in all psychosocial health indicators of reintegration, self-esteem, depressive symptoms, quality of life, PTSD, stigma and social support (Figure 1; Table 2). Psychosocial indicators were standardised to range 0–100; unstandardised results are presented in Table S2. Mean reintegration score at baseline was 33.2, increased substantially to 70.4 at 3 months post-surgery, gained an additional eight percentage points to 78.0 at 6 months and maintained through the end of the study. Self-esteem at baseline was mean 44.4, increased to 64.2 at 3 months and 72.4 at 6 months, and remained stable thereafter. Mean depressive symptoms also saw the greatest improvement from baseline to 3 months, (36.1–10.5), followed by a small decrease to 5.2 at 12 months. Over the 12 months, depressive symptomatology decreased from 85% to 7% (not shown).

We observed considerable elevations across all quality of life domains from baseline through 6 months post-surgery, but no further change through 12 months. The greatest mean increases from baseline to 6 months were for overall QOL (17.5–56.9) and the psychological health domain (24.4–65.7). Smaller increases, but still substantial, were observed for social relationships (38.2–49.5) and environment (41.0–58.9) domains. Ten participants (17%) screened positive for PTSD symptoms at baseline, reduced to one (2%) at 6 months and none at 12 months.

At baseline, the highest mean fistula-related stigma score was observed for the negative self-perception domain at 74.7 followed by the social isolation and verbal abuse domains (18.5 and 18.1, respectively). Mean score for fear of contagion was the lowest, at 6.7.


Physical Health

Fistula -related symptoms




Psychosocial Health

Reintegration, self-esteem, and depressive symptoms



Quality of life



Stigma




Figure 1 Trajectories of fistula-related physical symptoms and psychosocial health indicators across 12-month study follow-up.

Table 2 Physical and psychological health across 12-month study follow-up

	Baseline <i>n</i> = 60	3 months <i>n</i> = 59	6 months <i>n</i> = 55	9 months <i>n</i> = 55	12 months <i>n</i> = 58
Physical health					
General health					
Excellent	0 (0)	2 (3.4)	2 (3.6)	0 (0)	0 (0)
Very good	0 (0)	18 (31.5)	30 (54.6)	24 (43.6)	24 (41.4)
Good	0 (0)	15 (25.4)	7 (12.7)	9 (16.4)	16 (27.6)
Fair	11 (18.3)	17 (28.8)	12 (21.8)	18 (32.7)	17 (29.3)
Poor	49 (81.7)	7 (11.9)	4 (7.3)	4 (7.3)	1 (1.7)
Urinary incontinence					
Frequency of urinary incontinence	59 (98.3)	23 (38.9)	18 (32.7)	20 (36.4)	19 (32.8)
Never					
Once per week or less	1 (1.7)	36 (61)	37 (67.3)	35 (63.6)	39 (67.2)
Two or three times per week	0 (0)	1 (1.7)	1 (1.8)	3 (5.5)	5 (8.6)
Once per day	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Several times per day	2 (3.3)	4 (6.7)	1 (1.8)	2 (3.6)	4 (6.9)
All the time	12 (20)	7 (11.9)	6 (10.9)	7 (12.7)	4 (6.9)
Faecal incontinence					
All the time	45 (75)	11 (18.6)	10 (18.2)	8 (14.6)	6 (10.3)
Menstrual cycle*					
Regular	3 (5)	1 (1.7)	0 (0)	0 (0)	0 (0)
Irregular	<i>n</i> = 55 18 (32.7)	<i>n</i> = 54 29 (53.7)	<i>n</i> = 50 35 (70)	<i>n</i> = 50 33 (66.6)	<i>n</i> = 53 32 (60.4)
Amenorrhic	2 (3.6)	2 (3.7)	1 (2)	3 (6)	5 (9.4)
	35 (63.6)	23 (42.6)	14 (28)	14 (28)	16 (30.2)
Psychosocial health					
Reintegration	Mean (SD) 33.2 (20.4)	Mean (SD) 70.4 (27.1)	Mean (SD) 78.0 (23.7)	Mean (SD) 75.9 (25.7)	Mean (SD) 78.4 (19.2)
Self-esteem	44.4 (15.4)	64.2 (19.0)	72.4 (24.0)	69.9 (20.3)	70.7 (19.8)
Depressive symptoms	36.1 (16.9)	10.5 (13.2)	7.5 (9.6)	8.4 (11.8)	5.2 (7.7)
Quality of life					
Overall	17.5 (10.7)		56.9 (16.5)		54.2 (16.5)
Physical	47.9 (24.2)		81.4 (16.5)		75.3 (13.4)
Environment	41.0 (15.1)		58.9 (11.9)		60.2 (11.2)
Social relationships	38.2 (22.2)		49.5 (22.7)		52.6 (15.9)
Psychological	24.4 (15.1)		65.7 (21.6)		69.0 (17.4)
PTSD screen-positive <i>N</i> (%)	10 (16.7)		1 (1.8)		0 (0)
Stigma					
Negative self-perception	74.7 (36.2)	16.4 (32.5)	14.5 (29.4)	14.5 (32.9)	17.8 (33.6)
Social isolation	18.5 (27.2)	3.4 (10.1)	3.4 (13.3)	3.3 (10.3)	1.4 (6.1)
Verbal abuse	18.1 (26.5)	2.7 (7.8)	2.3 (7.2)	1.7 (6.4)	1.2 (3.8)
Fear of contagion	6.7 (18.4)	2.5 (8.7)	0.9 (5.0)	0.9 (5.0)	0 (0)
Social support	42.7 (10.0)	45.6 (12.3)	56.1 (20.5)	44.4 (11.6)	55.7 (19.4)

Mean (SD). Improvement is represented by increases in value for reintegration, self-esteem, and quality of life and decreases in depressive symptoms and stigma. Psychosocial measures are standardised to range 0–100.

*Among age <45 only.

Substantial declines were observed across all stigma domains during follow-up, with the largest improvements between baseline and 3 months. Mean stigma score improved by approximately 80% between baseline and 3 months for the domains of verbal abuse, social isolation and negative self-perception. Continued decline occurred through 12 months for verbal abuse (mean 1.2) and social isolation (mean 1.4). For negative self-perception, the decline continued through 6 months post-

surgery, but a small uptick occurred at 12 months (mean 17.8).

Psychosocial health among women with fistula-related physical symptoms

We assessed the relationship between urinary incontinence or other fistula-related physical symptoms and

Table 3 Unadjusted linear regression models of relationships between urinary incontinence or other fistula-related physical symptom and psychosocial health across study follow-up

	3 months		6 months		9 months		12 months		P
	B (95% CI)	P	B (95% CI)	P	B (95% CI)	P	B (95% CI)	P	
Reintegration measure	-24.1 (-38.8, -9.4)	0.0018	-18.2 (-30.2, -6.2)	0.0038	-22.6 (-35.6, -9.5)	0.001	-21.9 (-30.2, -13.5)	<0.0001	
Self-esteem	-19.3 (-29.3, -9.4)	0.0003	-9.1 (-19.8, 1.6)	0.095	-17.9 (-28.2, -7.6)	0.001	-21.2 (-30.1, -12.4)	<0.0001	
Depression	10.7 (3.4, 18)	0.0047	2.7 (-2.5, 7.9)	0.3071	8.3 (2.1, 14.5)	0.0098	7.8 (4.3, 11.3)	<0.0001	
Quality of life									
Overall			-15.8 (-23.8, -7.8)	0.0002			-22.5 (-28.9, -16.1)	<0.0001	
Physical			-17.6 (-25.3, -9.9)	<0.0001			-17.1 (-22.5, -11.7)	<0.0001	
Psychological			-17.9 (-28.7, -7.1)	0.0016			-19.1 (-26.8, -11.4)	<0.0001	
Social relationships			-13.1 (-25.1, -1.2)	0.0321			-13.1 (-20.7, -5.4)	0.0012	
Environment			-3.8 (-10.3, 2.7)	0.2443			-13.2 (-18, -8.4)	<0.0001	
Stigma									
Verbal abuse	3.7 (-0.8, 8.2)	0.106	0.3 (-3.6, 4.3)	0.8649	2.2 (-1.4, 5.7)	0.2236	2.3 (0.1, 4.5)	0.0415	
Negative self-perception	22.5 (4.2, 40.8)	0.0169	22.1 (7.1, 37.1)	0.0046	23.5 (6.2, 40.8)	0.0086	34 (18.8, 49.1)	<0.0001	
Social isolation	4.9 (-1, 10.8)	0.0989	2.4 (-4.9, 9.7)	0.5122	5.4 (-0.2, 10.9)	0.0568	4.2 (-0.7, 9.1)	0.091	
Fear of contagion	3.5 (-1.5, 8.5)	0.1706	-0.9 (-3.6, 1.9)	0.5352	1.5 (-1.3, 4.2)	0.2923	0.6 (-0.6, 1.7)	0.3048	

Improvement is represented by increases in value for reintegration, self-esteem, and quality of life and decreases in depressive symptoms and stigma. At each time point, the values presented reflect the difference between those currently reporting symptoms *vs.* those reporting no symptoms. Psychosocial measures are standardised to range 0–100.

indicators of women's psychosocial health (Table 3). Across study follow-up, self-report of physical symptoms largely correlated with significantly lower psychosocial health. Mean reintegration score differed by about 20 points at each time point comparing women reporting physical symptoms to those reporting no symptoms. At 12 months, women reporting physical symptoms had a mean reintegration score 21.9 points lower than women reporting no physical symptoms (95% CI: -30.2, -13.5). Self-esteem among women reporting physical symptoms was also significantly lower: this difference ranged in magnitude from mean 19.3 points lower at 3 months (95% CI -29.3, -9.4) to 21.2 points lower at 12 months (95% CI -30.1, -12.4). Women reporting physical symptoms had significantly higher mean depression at 3, 9 and 12 months by about eight points. At 12 months, women reporting physical symptoms scored on average 7.8 points higher (95% CI 4.3, 11.3) on the standardised depression measure compared to those with no symptoms. Physical symptoms were associated with substantial differences in quality of life domains as well. At 12 months, we observed an approximately 20-point lower value in overall quality of life (-22.5, 95% CI -28.9, -16.1) and in the physical (-17.1, 95% CI -22.5, -11.7) and psychological domains (-19.1, 95% CI -26.8, -11.4) for women reporting physical symptoms compared to those reporting no symptoms. Slightly lower differences were observed for quality of life domains of social relationships (-13.1, 95% CI -20.7, -5.4) and environment (-13.2, 95% CI -18.0, -8.4). Differences in stigma by the report of physical symptoms were less consistent. Across all time points, large and statistically significant differences in the stigma domain of negative self-perception were observed, with a mean 22.5 at baseline (95% CI 4.2–40.8) and rising to 34.0 (95% CI 18.8–49.1) at 12 months. Smaller and largely non-significant differences were identified across symptom categories for the stigma domains social isolation, verbal abuse and fear of contagion. The reduction in difference for social isolation by 12 months was less than that observed for the other domains.

Discussion

Our study is the first to prospectively assess trajectories of physical and psychosocial recovery in the first year following surgery for obstetric fistula. In this longitudinal cohort of women in Uganda, women experienced dramatic improvements in physical and psychosocial health following surgery. The most substantial gains were typically observed by 3 months with continued improvement through 6 months, after which little change was

observed. Further, we quantified the large disparity in psychosocial health between women with persistent physical symptoms compared to those without, identifying significant differences across most indicators.

The most common persistent physical symptoms at 12 months post-surgery were urinary incontinence (32.8%) and weakness (17.2%). While the 12-month values represent a substantial reduction from baseline, the proportion of women with persistent symptoms is nonetheless concerning. Our findings on urinary incontinence were on the higher end of those reported in other studies reporting persistent incontinence rates of 16–35% [13, 18, 32, 53–55]. Our identification of non-incontinence-related persistent symptoms is also not unique; in a qualitative study of women repaired for fistula in Ethiopia, Donnelley *et al.* [28] reported that 40% of women reported physical health problems following surgery irrespective of their repair status. The substantial and significant differences in level of reintegration comparing women with persistent physical symptoms and those without support additional intervention to resolve physical symptoms in order to support women's recovery or more targeted psychosocial support and reintegration services to ensure that those coping with physical or psychosocial challenges are appropriately supported.

Our findings demonstrating dramatic improvement in quality of life following surgery are consistent with other reports [20, 24, 31] although several interesting differences in mean value and changes over time across setting are notable in two studies in India and Nigeria also using the WHO QOL-BREF that followed women through 6 months (Nigeria) and median 26 months (India) after surgery [56, 57]. The lowest pre-operative QOL score for all domains except social relationships and the largest improvements from the pre- to the post-operative period were in India. In India and Nigeria, significant improvements were noted in the physical, psychosocial and social domains but no change was observed for the environment domain. The largest improvements in Uganda and India were observed in the physical domain. Further comparison with population-based samples of healthy women of reproductive age would help to place these data into context. Additional research on the contextual factors responsible for the differences identified in overall experience and change across time could help inform efforts to improve recovery trajectories across various cultural contexts.

We found substantial reductions across the dimensions of fistula-related stigma for the 12-month period, with the steepest declines observed from baseline through 3 months. In this cohort, negative self-perception was the most severe type of stigma experienced across all time

points. Compared to negative self-perception, the relatively lower report of stigma domains representing enacted stigma (verbal abuse, social isolation and fear of contagion) may be partially explained by nearly half of our cohort achieving surgery within 3 months of fistula development. Our finding of persistent stigma is consistent with one qualitative study of Ethiopian women's post-surgical quality of life, where persistent internalised and enacted stigma were reported at mean 1.5 years following surgery irrespective of surgical outcome [28]. Literature from other stigmatised conditions describes a complex relationship between perceived community stigma and internalised stigma [58] and supports a strong relationship between internalised stigma and depression and other poor health outcomes [59–61], highlighting the need for psychosocial intervention at multiple levels to reduce stigma experiences and associated consequences [62, 63].

The substantial differences observed in psychosocial outcomes comparing women with physical symptoms to those without highlight the need for identification of strategies to treat or better manage post-surgical fistula-related symptoms. Our results suggest that the greatest overall physical health benefits occur during the first 6 months following surgery, with concurrent psychosocial health improvement. The stagnation in physical and psychosocial health indicators that we observed at about 6 months is consistent with qualitative work conducted Kenya by Khisa *et al.* [64]. Khisa describes a 'miracle phase' in the immediate post-surgical period among women with successful repair. While some of the stagnation in trend may be due to floor and ceiling effects, our results may be also be influenced by individual experiences not explored here including fistula-related characteristics or experiences, sexual relationships, concerns around achieving or preventing pregnancy and exposure to violence. Other factors, such as poverty, lack of empowerment and lack of social support, may also structure health status, and thus, our findings may represent physical and psychosocial health status comparable to similar women of reproductive age in this setting who did not develop fistula. Subsequent work should explore the contributions of other factors, and development of interventions to improve recovery will benefit from the inclusion of theory-based conceptual models for recovery from mental health diagnoses suggest areas for counselling foci that are particularly salient for women recovering from fistula, based on our findings here and other literature [65].

We recognise several limitations to this study. Our sample size was based on feasibility; thus, our results should be validated in larger samples. Despite our small

sample, our rates of participation were high, and our cohort was representative of women accessing fistula repair at Mulago Hospital during the recruitment period. Both similarities and important differences in women's experiences of fistula and recovery are observed in the literature, and thus, a nuanced assessment of the contextual factors contributing to these differences across countries will help establish the generalisability of our results. In particular, many women in our cohort rapidly accessed surgery and nearly thirty per cent of infants survived the labour resulting in the fistula, suggesting improved care access for this cohort compared to some other studies. Our sample included participants with childbirth-related fistula due to pressure necrosis from obstructed labour and iatrogenic aetiologies; however, designation of likely iatrogenic among this sample was retrospectively assigned using medical record criteria [36]. Our findings can be considered generally representative of women with obstetric fistula, but further research exploring differences in recovery trajectories by aetiology is of interest, particularly as iatrogenic female genital fistula appears to be increasing [66]. Our study is both strengthened and limited by the length of follow-up; our focus on the 12 months following surgery expands the length of follow-up conducted in many other studies, yet may be inadequate to understand the rates and consequences of fistula recurrence and adverse perinatal outcome over a woman's reproductive life, both of which have been found to be high among women receiving surgical repair for obstetric fistula [67].

Reducing the impact of fistula-related maternal morbidity on the health of women requires their engagement in the full continuum of fistula-related management. Prevention is key, and improving women's timely access to surgical repair of fistula is critical. Yet limiting attention to women recovering from fistula in the immediate post-surgical period is inadequate. Intervention programming to support the spectrum of women's physical and psychosocial needs during their recovery from fistula may result in improved outcomes after repair. Efforts must also be made to maintain women's engagement with health systems and to improve the responsiveness of health systems to women's needs following surgery to protect maternal health and well-being.

Acknowledgements

The authors would like to recognise our study participants for their commitment to this study and thank Erin Mielke and Mary Ellen Stanton for their comments and Amy J. Markowitz, JD, for her editorial support. Luganda translations for WHO QOL-BREF and the

Hopkins Symptom Checklist were created by The School of International Development, University of East Anglia, UK; and The Social Science Programme, Medical Research Council/UVRI Uganda Research Unit on AIDS, Entebbe, Uganda. Funding for open source publication was provided by the Fistula Care *Plus* Project, administered by EngenderHealth (cooperative agreement AID-OAA-A14-00013), and the opinions expressed herein are those of the authors and do not necessarily reflect the views of USAID, or the United States Government.

Funding

The study was funded by the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the United States National Institutes of Health (NICHD; Project number R21HD075008), the Fistula Foundation and from the United States Agency for International Development (USAID) via the Fistula Care *Plus* Project, administered by EngenderHealth (cooperative agreement AID-OAA-A14-00013). The opinions expressed are those of the authors and do not necessarily reflect the views of USAID, or the United States Government. Continuing analytical work was funded by NICHD (Project number K99HD086232).

Ethics approval

The study protocol was approved by the Makerere University College of Health Sciences, School of Medicine Research and Ethics Committee (#2014-052, Apr 2014), the Uganda National Council for Science and Technology (#154/212/01, June 2015), and the University of California, San Francisco Human Research Protection Program, Committee on Human Research (#12-09573, Oct 2012).

References

- Arrowsmith S, Hamlin EC, Wall LL. Obstructed labor injury complex: obstetric fistula formation and the multifaceted morbidity of maternal birth trauma in the developing world. *Obstet Gynecol Surv* 1996; **51**: 568–574.
- Ahmed S, Holtz SA. Social and economic consequences of obstetric fistula: life changed forever? *Int J Gynaecol Obstet* 2007; **99**(Suppl 1): S10–S15.
- Browning A, Member B. Women with obstetric fistula in Ethiopia: a 6-month follow-up after surgical treatment. *BJOG* 2008; **115**: 1564–1569.
- Roush KM. Social implications of obstetric fistula: an integrative review. *J Midwifery Womens Health* 2009; **54**: e21–e33.
- Turan JM, Johnson K, Polan ML. Experiences of women seeking medical care for obstetric fistula in Eritrea: implications for prevention, treatment, and social reintegration. *Glob Public Health* 2007; **2**: 64–77.
- Siddle K, Mwambingu S, Malinga T, Fiander A. Psychosocial impact of obstetric fistula in women presenting for surgical care in Tanzania. *Int Urogynecol J* 2013; **24**: 1215–1220.
- Ashford L. *Obstetric Fistula: Cause of Stigma and Disability in Less Developed Countries*. Population Reference Bureau: Washington, D.C., 2002.
- Bangser M. Obstetric fistula and stigma. *Lancet* 2006; **367**: 535–536.
- Gebresilase YT. A qualitative study of the experience of obstetric fistula survivors in Addis Ababa, Ethiopia. *Int J Womens Health* 2014; **6**: 1033–1043.
- Ojengbede OA, Baba Y, Morhason-Bello IO *et al.* Group psychological therapy in obstetric fistula care: a complementary recipe for the accompanying mental ill health morbidities? *Afr J Reprod Health* 2014; **18**: 155–159.
- Zelege BM, Ayele TA, Woldetsadik MA, Bisetegn TA, Adane AA. Depression among women with obstetric fistula, and pelvic organ prolapse in northwest Ethiopia. *BMC Psychiatry* 2013; **13**: 236.
- Weston K, Mutiso S, Mwangi JW, Qureshi Z, Beard J, Venkat P. Depression among women with obstetric fistula in Kenya. *Int J Gynaecol Obstet* 2011; **115**: 31–33.
- Wall LL. Obstetric vesicovaginal fistula as an international public-health problem. *Lancet* 2006; **368**: 1201–1209.
- WHO. *Obstetric Fistula: Guiding Principles for Clinical Management and Programme Development*. WHO: Geneva, 2006.
- Adler AJ, Ronsmans C, Calvert C, Filippi V. Estimating the prevalence of obstetric fistula: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2013; **13**: 246.
- Kayondo M, Wasswa S, Kabakyenga J *et al.* Predictors and outcome of surgical repair of obstetric fistula at a regional referral hospital, Mbarara, western Uganda. *BMC Urol* 2011; **11**: 23.
- de Bernis L. Obstetric fistula: guiding principles for clinical management and programme development, a new WHO guideline. *Int J Gynaecol Obstet* 2007; **99**(Suppl 1): S117–S121.
- Delamou A, Diallo M, Beavogui AH *et al.* Good clinical outcomes from a 7-year holistic programme of fistula repair in Guinea. *Trop Med Int Health* 2015; **20**: 813–819.
- Arrowsmith SD, Ruminjo J, Landry EG. Current practices in treatment of female genital fistula: a cross sectional study. *BMC Pregnancy Childbirth*. 2010; **10**: 73.
- Browning A. Risk factors for developing residual urinary incontinence after obstetric fistula repair. *Br J Obstet Gynaecol* 2006; **113**: 482–485.
- Pope R, Bangser M, Requejo JH. Restoring dignity: social reintegration after obstetric fistula repair in Ukerewe, Tanzania. *Glob Public Health* 2011; **6**: 859–873.
- Wilson SM, Sikkema KJ, Watt MH, Masenga GG, Mosha MV. Psychological symptoms and social functioning

A. M. El Ayadi *et al.* **Physical and psychosocial health after fistula surgery**

- following repair of obstetric fistula in a low-income setting. *Matern Child Health J* 2016; **20**: 941–945.
23. Browning A, Menber B. Women with obstetric fistula in Ethiopia: a 6 month follow up after surgical treatment. *Br J Obstet Gynaecol*. 2008; **15**: 1564–1569.
 24. Wilson SM, Sikkema KJ, Watt MH, Masenga GG. Psychological symptoms among obstetric fistula patients compared to gynecology outpatients in Tanzania. *Int J Behav Med* 2015; **22**: 605–613.
 25. Landry E, Frajzyngier V, Ruminjo J *et al.* Profiles and experiences of women undergoing genital fistula repair: findings from five countries. *Glob Public Health* 2013; **8**: 926–942.
 26. Women's Dignity Project, EngenderHealth. *Sharing the Burden: Ugandan Women Speak about Obstetric Fistula*. Women's Dignity Project and EngenderHealth: Dar es Salaam, Tanzania, 2007.
 27. Bangser M. Strengthening public health priority-setting through research on fistula, maternal health, and health inequities. *Int J Gynaecol Obstet* 2007; **99**(Suppl 1): S16–S20.
 28. Donnelly K, Oliveras E, Tilahun Y, Belachew M, Asnake M. Quality of life of Ethiopian women after fistula repair: implications on rehabilitation and social reintegration policy and programming. *Cult Health Sex* 2015; **17**: 150–164.
 29. Anzaku SA, Lengmang SJ, Mikah S, Shephard SN, Edem BE. Sexual activity among Nigerian women following successful obstetric fistula repair. *Int J Gynaecol Obstet* 2017; **137**: 67–71.
 30. Khisa AM, Nyamongo IK. Still living with fistula: an exploratory study of the experience of women with obstetric fistula following corrective surgery in West Pokot, Kenya. *Reprod Health Matters* 2012; **20**: 59–66.
 31. El-Gazzaz G, Hull TL, Mignanelli E, Hammel J, Gurland B, Zutshi M. Obstetric and cryptoglandular rectovaginal fistulas: long-term surgical outcome; quality of life; and sexual function. *J Gastrointest Surg* 2010; **14**: 1758–1763.
 32. Nielsen HS, Lindberg L, Nygaard U *et al.* A community-based long-term follow up of women undergoing obstetric fistula repair in rural Ethiopia. *BJOG* 2009; **116**: 1258–1264.
 33. Wall LL, Arrowsmith SD, Briggs ND, Lassey A. Urinary incontinence in the developing world: the obstetric fistula. In: Abrams P, Cardozo L, Khoury S, Wein A (eds). *Incontinence*. Health Publication Ltd: Plymouth, UK, 2002.
 34. Wall LL. Dead mothers and injured wives: the social context of maternal morbidity and mortality among the Hausa of northern Nigeria. *Stud Fam Plann* 1998; **29**: 341–359.
 35. Byamugisha J, El Ayadi A, Obore S *et al.* Beyond repair – family and community reintegration after obstetric fistula surgery: study protocol. *Reprod Health* 2015; **12**: 115.
 36. Raassen TJ, Ngongo CJ, Mahendeka MM. Iatrogenic genitourinary fistula: an 18-year retrospective review of 805 injuries. *Int Urogynecol J* 2014; **25**: 1699–1706.
 37. Lorig K, Steward A, Ritter P, Gonzalez V, Laurent D, Lynch J. *Outcome Measures for Health Education and Other Health Care Interventions*. Sage Publications: Thousand Oaks, CA, 1996.
 38. Klovning A, Avery K, Sandvik H, Hunskaar S. Comparison of two questionnaires for assessing the severity of urinary incontinence: the ICIQ-UI SF versus the incontinence severity index. *Neurourol Urodyn* 2009; **28**: 411–415.
 39. El Ayadi A, Byamugisha J, Obore S *et al.* Development and preliminary validation of a post-fistula repair reintegration instrument among Ugandan women. *Reprod Health J* 2017; **14**: 109.
 40. Bolton P, Wilk CM, Ndogoni L. Assessment of depression prevalence in rural Uganda using symptom and function criteria. *Soc Psychiatry Psychiatr Epidemiol* 2004; **39**: 442–447.
 41. Derogatis LR, Lipman RS, Rickels K, Uhlenhuth EH, Covi L. The Hopkins Symptom Checklist (HSCL): a self-report symptom inventory. *Behav Sci* 1974; **19**: 1–15.
 42. Martin F, Russell S, Seeley J. The WHOQOL BREF questionnaire in Luganda: Validation with a sample including people living with HIV in Uganda. Working Paper 46. School of International Development, University of East Anglia: Norwich, UK, 2013.
 43. Webster J, Nicholas C, Velacott C, Cridland N, Fawcett L. Validation of the WHOQOL-BREF among women following childbirth. *Aust N Z J Obstet Gynaecol* 2010; **50**: 132–137.
 44. Rosenberg M. *Society and the Adolescent Self-image*. Princeton University Press: Princeton, NJ, 1965.
 45. Holzemer WL, Uys LR, Chirwa ML *et al.* Validation of the HIV/AIDS stigma instrument – PLWA (HASI-P). *AIDS Care* 2007; **19**: 1002–1012.
 46. Prins A, Ouimette P, Kimerling R *et al.* The primary care PTSD screen (PC-PTSD): development and operating characteristics. *Primary Care Psychiatry* 2003; **9**: 9–14.
 47. Zimet GD, Powell SS, Farley GK, Werkman S, Berkoff KA. Psychometric characteristics of the Multidimensional Scale of Perceived Social Support. *J Pers Assess* 1990; **55**: 610–617.
 48. Nakigudde J, Musisi S, Ehnvall A, Airaksinen E, Agren H. Adaptation of the multidimensional scale of perceived social support in a Ugandan setting. *Afr Health Sci* 2009; **9**(Suppl 1): S35–S41.
 49. Bolton P, Bass J, Neugebauer R *et al.* Group interpersonal psychotherapy for depression in rural Uganda: a randomized controlled trial. *JAMA* 2003; **289**: 3117–3124.
 50. Kaida A, Matthews LT, Ashaba S *et al.* Depression during pregnancy and the postpartum among HIV-infected women on antiretroviral therapy in Uganda. *J Acquir Immune Defic Syndr* 2014; **67**(Suppl 4): S179–S187.
 51. Psaros C, Haberer JE, Boum Y 2nd *et al.* The factor structure and presentation of depression among HIV-positive adults in Uganda. *AIDS Behav* 2015; **19**: 27–33.
 52. Ashaba S, Kakuhihike B, Vorechovska D *et al.* Reliability, validity, and factor structure of the Hopkins Symptom Checklist-25: population-based study of persons living with

A. M. El Ayadi *et al.* **Physical and psychosocial health after fistula surgery**

- HIV in rural Uganda. *AIDS Behav* 2018; **22**: 1467–1474.
53. Semere L, Nour NM. Obstetric fistula: living with incontinence and shame. *Rev Obstet Gynecol* 2008; **1**: 193–197.
54. Lewis A, Kaufman MR, Wolter CE *et al.* Genitourinary fistula experience in Sierra Leone: review of 505 cases. *J Urol* 2009; **181**: 1725–1731.
55. Nardos R, Browning A, Chen CC. Risk factors that predict failure after vaginal repair of obstetric vesicovaginal fistulae. *Am J Obstet Gynecol* 2009; **200**(578): e1–e4.
56. Singh V, Jhanwar A, Mehrotra S, Paul S, Sinha RJ. A comparison of quality of life before and after successful repair of genitourinary fistula: Is there improvement across all the domains of WHOQOL-BREF questionnaire? *Afr J Urol* 2015; **21**: 230–234.
57. Umoiyoho AJ, Inyang-Etoh EC, Abah GM, Abasiattai AM, Akaiso OE. Quality of life following successful repair of vesicovaginal fistula in Nigeria. *Rural Remote Health* 2011; **11**: 1734.
58. Turan B, Budhwani H, Fazeli PL *et al.* How does stigma affect people living with HIV? The mediating roles of internalized and anticipated HIV stigma in the effects of perceived community stigma on health and psychosocial outcomes. *AIDS Behav* 2017; **21**: 283–291.
59. Turan B, Hatcher AM, Weiser SD, Johnson MO, Rice WS, Turan JM. Framing mechanisms linking HIV-related stigma, adherence to treatment, and health outcomes. *Am J Public Health* 2017; **107**: 863–869.
60. Turan B, Smith W, Cohen MH *et al.* Mechanisms for the negative effects of internalized HIV-related stigma on antiretroviral therapy adherence in women: the mediating roles of social isolation and depression. *J Acquir Immune Defic Syndr* 2016; **72**: 198–205.
61. Sweeney SM, Venable PA. The association of HIV-related stigma to HIV medication adherence: a systematic review and synthesis of the literature. *AIDS Behav* 2016; **20**: 29–50.
62. Stangl AL, Lloyd JK, Brady LM, Holland CE, Baral S. A systematic review of interventions to reduce HIV-related stigma and discrimination from 2002 to 2013: how far have we come? *J Int AIDS Soc* 2013; **16**(3 Suppl 2): 18734.
63. Takada S, Weiser SD, Kumbakumba E *et al.* The dynamic relationship between social support and HIV-related stigma in rural Uganda. *Ann Behav Med* 2014; **48**: 26–37.
64. Khisa W, Wakasiaka S, McGowan L, Campbell M, Laverder T. Understanding the lived experience of women before and after fistula repair: a qualitative study in Kenya. *BJOG* 2017; **124**: 503–510.
65. Leamy M, Bird V, Le Boutillier C, Williams J, Slade M. Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *Br J Psychiatry* 2011; **199**: 445–452.
66. Wright J, Ayenachew F, Ballard KD. The changing face of obstetric fistula surgery in Ethiopia. *Int J Womens Health* 2016; **8**: 243–248.
67. Delamou A, Delvaux T, El Ayadi AM *et al.* Fistula recurrence, pregnancy, and childbirth following successful closure of female genital fistula in Guinea: a longitudinal study. *Lancet Glob Health*. 2017; **5**: e1152–e1160.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Description of physical and psychosocial health measures.

Table S2. Psychosocial health indicators across study 12-month study follow-up, Unstandardised.

Corresponding author Alison M. El Ayadi, Department of Obstetrics, Gynecology and Reproductive Sciences, Bixby Center for Global Reproductive Health, 550 16th Street, UCSF Box 1224, San Francisco, CA 94158, USA. Tel.: +1-415-659-8367; E-mail: alison.elayadi@ucsf.edu