

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/342158462>

Preliminary Antidiabetic Potential of Ugandan- Matooke (Musa paradisiaca) Peels

Article · June 2020

DOI: 10.9734/IJBCRR/2020/v29i530190

CITATIONS

0

READS

224

5 authors, including:



Adaku Iwueke

University of Agriculture and Environmental Sciences Umuagwo Nigeria

10 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)



John Nnamdi Ejekwumadu

Teesside University

8 PUBLICATIONS 6 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Probiotic potentials of lactic acid bacteria isolated from amakamo, a fermented milk from Uganda. [View project](#)



Mental health and Wellness [View project](#)



Preliminary Antidiabetic Potential of Ugandan-Matooke (*Musa paradisiaca*) Peels

Adaku Vivien Iwueke^{1*#}, Nnamdi John Ejekwumadu^{1##}, Roland Kiiza²,
Kenneth Iceland Kasozi³ and Swafia Khambari¹

¹Department of Biochemistry, Kampala International University, Western Campus Ishaka-Bushenyi, Uganda.

²Department of Pharmacology, Kampala International University, Western Campus Ishaka-Bushenyi, Uganda.

³Department of Physiology, Kampala International University, Western Campus Ishaka-Bushenyi, Uganda.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AVI, NJE conceptualized and designed the study. Authors RK, NJE and SK carried out the laboratory experiments. Authors RK, KIK did the literature search, data analysis and drafting of the manuscript. Authors AVI, NJE critically reviewed the concept, study design, literature, data analysis and edited the manuscript. All the authors read and consented to the final manuscript.

Article Information

DOI: 10.9734/IJBCRR/2020/v29i530190

Editor(s):

(1) Prof. Halit Demir, Yil University, Turkey.

Reviewers:

(1) Bongani Alphonse Nkala, Pearson Institute of Higher Education, South Africa.

(2) Augustine Ikhueoya Airaodion, Federal University of Technology Owerri, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/57725>

Original Research Article

Received 01 April 2020

Accepted 06 June 2020

Published 13 June 2020

ABSTRACT

Background: Diabetes mellitus is a metabolic disease characterized by hyperglycaemia over a prolonged period. In Uganda, unripe *Musa paradisiaca* (banana) is a staple food which is deskinning before cooking. In West Africa, however, the unpeeled banana is cooked for diabetics.

Aim: The objective of the study was to assess the hypoglycemic and body weight changes in experimental diabetic Wistar rats treated with green banana peel extracts.

Methodology: Out of the thirty-nine male rats used for the study, nine were used for the evaluation of possible oral acute toxicity. The remaining thirty were divided into six groups of 5 rats each. Experimental diabetes mellitus was induced by intravenous injection of alloxan (65 mg/kg body

*Corresponding author: E-mail: drvivien.89@gmail.com, aiwueke@pums.edu.ng;

[#]Department of Biochemistry, PAMO University of Medical Sciences, Port-Harcourt, Nigeria.

^{##}Imo State Polytechnic, Umuagwo, Nigeria.

weight) the alloxan and treated with insulin, distilled water and graded doses of the extract (1000 mg/kg, 2000 mg/kg and 4000 mg/kg body weight) for two weeks.

The weight and blood glucose levels were measured before and after induction and prior to administration of treatment dosages. Data were analyzed using SPSS Version 20. ANOVA and spearman's rank correlation tests determined significant changes in values at 95% confidence interval.

Results: The rats showed no signs of toxicity nor death even at a dose of 10,000 mg/kg body weight. Phytochemical screening revealed saponins, tannins, phenols, flavonoids, cardiac glycoside, alkaloids, steroids and terpenoids. This study demonstrates that *Musa paradisiaca* peels significantly attenuated blood glucose levels ($P < 0.005$) and regulated body weights at doses (≥ 2000 mg/kg) which are essential parameters in the management of diabetes mellitus.

Conclusion: In line with the findings, unripe banana peels are anti-diabetic; unripe bananas therefore, should be cooked with the skins to derive its established anti-diabetic benefits.

Keywords: *Diabetes mellitus; Musa paradisiaca; matooke peels; medicinal plants; phytochemicals.*

1. INTRODUCTION

About 442 million people worldwide have diabetes mellitus WHO [1]. Africa is undergoing a rapid demographic transition with the burden of non-communicable diseases, especially diabetes, that will further overload the health care systems already challenged by HIV/AIDS and Malaria [2]. All these are due to rapid urbanization and the attendant changes in lifestyle, rapidly decreasing physical activity, nutrition and ageing of the population [3]. Diabetes mellitus (DM), also known as diabetes, is a group of metabolic diseases which manifest as high blood sugar levels [3]. DM presents with inadequate insulin production, or body cells do not respond appropriately to the insulin, or both, affecting multiple systems in the process [4]. Symptoms of high blood glucose include frequent urination, increased thirst and increased hunger [5]. If left untreated, diabetes can lead to many complications [5]. Acute complications include diabetic ketoacidosis and non-ketotic hyperosmolar coma [5]. Grave long-term complications include cardiovascular disease, stroke, chronic kidney failure, foot ulcers and damage to the eyes [6,7].

About 25 million African adults are estimated to have diabetes [1]. In Uganda, DM prevalence varies between 90.9% [8] and 31% [9]. Bearing in mind that there is no completely effective treatment available for DM, majority of communities in sub-Saharan Africa have adopted the use of ethno medicinal plants for the management of clinical complications associated with the disease [10]. Studies show that members of the *Musaceae* family have significant potentials in managing diabetes [11]. The family also alleviates the toxic effects of

carbon tetrachloride in animals as well as a profound *in vitro* and *in vivo* anthelmintic activity [12,13]. Different parts of the banana plant have been demonstrated to possess detectable antidiabetic activity [14]. These have included fruits, leaves, roots and even flowers [14,15], leading to the speculation about the banana peels. *Matooke* is a Ugandan staple, prepared by peeling the unripe bananas before cooking and mashing the soft bananas while hot.

2. MATERIALS AND METHODS

2.1 Collection of Plant Material and the Extraction Method

The bananas were collected from a single plant of the common East African Highland Cultivar of *Musa paradisiaca* in a farm at Ishaka Bushenyi, Uganda. Preparation of unripe green banana peel extract was done according to previous study [16] with minor modifications. After harvesting the green banana, it was peeled and the peels were air dried on a laboratory bench for 1 week. The dried banana peels (100 grams) in 1000 ml of water were boiled for 20 minutes and blended for five minutes on cooling. The resultant solution was then filtered with sterile wire gauze to remove the fibers. Subsequent filtration was carried out using Watman No.1 filter paper. The extract was poured into a weighed glass beaker and evaporated at 100°C in a water bath to obtain a semi-solid crude extract which was further dried in an oven at 60°C.

2.2 Animals

The study was carried out in the Pharmacology and Biochemistry laboratories of the School of Health Sciences, Kampala International

University, Western Campus Ishaka. The animals were obtained from the animal housing facility of the School of Pharmacy, Kampala International University. They were fed standard animal pellets and given water *ad libitum*.

2.2.1 Sampling technique

The experimental animals were acclimatized for seven days before the commencement of the experiment and randomly placed in groups and appropriate labels were used to identify the animals.

2.2.2 Inclusion/exclusion criteria

Mature male Wistar rats weighing above 100 g were used in the study; while rats weighing less than 100 g were excluded.

2.3 Phytochemical Analysis

The phytochemical analysis of the extract of *Musa paradisiaca* peels was carried out to identify the phytochemical constituents present in the extract. Tests were carried out using standard methods [17].

2.4 Determination of Acute Toxicity

The acute toxicity of the aqueous extract of the green banana was determined by using 9 Wistar rats as described by [18]. The animals were fasted for 24 hours prior to experiment. Animals were weighed and labelled with different body marks. Nine animals were grouped into three and a different dose was administered to each group.

2000 mg/kg, 4000 mg/kg and 10000 mg/kg doses were used in the study.

The corresponding volume to be administered was calculated using the formula:

$$\frac{A \times B}{C \times 1000}$$

Where,

A = Body Weight; B = Dose of administration; C = Concentration

The animals were observed for signs of toxicity after 3 hours and 24 hours respectively.

2.5 Induction of Experimental Diabetes

Rats were weighed, marked and fasted for 12 hours. Afterwards, diabetes was induced by a

single intraperitoneal administration of alloxan monohydrate (65 mg/kg) with 4% saline solution (an average of 0.90 mL per test animal). Before the administration of alloxan, a base line glycaemia was recorded from blood collected by tail vein puncture of each rat.

After 3 days, the rats' blood glucose levels were determined by testing them with diabetes test strips and glucometer (OnCall®). Rats with blood glucose above 200 mg/dl (7.0 mMols) were considered diabetic and selected for the study. One group of the rats to which alloxan was not administered served as normal control.

2.6 Treatment

The animals were randomized into 6 groups, each group consisting of 5 test animals [18] as follows;

Group I: Non-diabetic rats received distilled water (DW) (10 ml/kg) served as normal control group.

Group II: Diabetic rats received DW (10 ml/kg) served as diabetic control group.

Group III: Diabetic rats received aqueous extract of banana peel at a dose of 1000 mg/kg.

Group IV: Diabetic rats received aqueous extract of banana peel at a dose of 2000 mg/kg.

Group V: Diabetic rats received aqueous extract of banana peel at the dose of 4000 mg/kg.

Group VI: Diabetic rats received insulin at dose of 10 U/kg as a control reference.

Oral administration of distilled water (DW) and peel extract was done daily before feeding the animals. Insulin treatment (10 U) administered intra-muscular was used as reference point. The final blood sugar was measured after two weeks.

2.7 Data Analysis

Data were transferred into Microsoft Excel spreadsheet 2016, analysed and information expressed as mean \pm SD at 95% confidence interval. Effects of treatment on blood glucose and body weight was analysed using SPSS Version 20 and the results were expressed as mean \pm SD. Comparisons were carried by analysis of variance (ANOVA) to determine dosage groups that are important in causing changes in Type 1 Diabetes Mellitus (T1DM)

Wistar rats. Significance was set at $P \leq 0.05$. Correlation analysis was conducted on body weight and blood glucose levels to determine the level of relationship and a spearman's correlation test was done at 95% confidence interval.

3. RESULTS AND OBSERVATIONS

Phytochemical analysis showed the presence of important secondary metabolites (Table 1).

Table 1. Phytochemicals present in the aqueous extract of green banana peels

Saponnins	+
Tannins	+
Phenols	+
Flavonoids	+
Cardiac glycosides	+
Alkaloids	+
Steroids	Trace
Terpenoids	+
Phlobatannins	-
Antraquinone	-

3.1 Body Weights and Serum Blood Glucose Levels in Treatment Groups

The results showed that the mean body weight was 143.48-157.27 g in the extract treatment groups. Insulin treated group had a mean body weight of 152.96 ± 5.99 g. The mean serum blood glucose levels in the insulin treated group was 11.25 ± 1.18 mMol/dl while in the extract treatment group, they were in the range of 10.12-10.58 mMol/dl at 95% confidence interval as shown in Table 2.

4. DISCUSSION

The acute toxicity study showed no signs of toxicity such as salivation, diarrhoea, depression,

stimulation, vomiting, and coma, and there was no mortality observed up to a dose of 10,000 mg/kg. The extract proved to be safe for short term administration. However, a chronic toxicity study will be appropriate to see any cumulative tissue or organ impairment.

From the study, in the diabetic groups the mean body weights are above 140 kg while the mean blood glucose levels are above 11 mMol/dl. Control of hyperglycemia in diabetes sometimes results in weight gain and hypoglycemia, which is in agreement with a previous study [19]. This would be the primary underlying factor as to why there is a general global weight gain amongst DM patients [20]. The study further showed moderate weight gains in the treatment groups on the extract, which confirms the previously held opinion that vegetable diets protect against weight gain in DM [21]. Showing that understanding the physiology of weight gain in the management of Type 1 Diabetes Mellitus (T1DM) is crucial to gain the desired therapeutic outcome since weight gain is strongly associated with increased risk for DM [22]. Moreover, excessive weight gain is a major contributory factor to poor glycemic control in DM, thus showing the need to have therapeutic options which effectively control it [23].

The study also demonstrated that serum blood glucose levels are in the range of 11.12-12.57 mMol/dl in groups treated with banana peels. Similar results were shown in the Insulin treatment group (11.25 ± 10.2 mMol/dl), justifying the hypoglycemic effect of *Musa paradisiaca* peels. A previous study has demonstrated the hypoglycemic effect of the flesh of the green fruit [15]. This study has been able to demonstrate that *Musa paradisiaca* peels have a potent hypoglycemic effect at relatively higher doses. Since, relatively, no effect had

Table 2. Body weight and glucose levels in treatment groups

Treatment groups	N	Body weight				Serum blood glucose levels(mMol/L)			
		Mean \pm	SD	95% CI		Mean \pm	SD	95% CI	
				LL	UL			LL	UL
Insulin (10 U)	5	152.96	5.99	151.96	153.96	11.25	1.18	10.252	12.248
Control	5	150.42	16.78	149.33	151.51	4.41	0.44	3.317	5.503
1000 mg/kg	5	147.54	41.86	146.54	148.54	11.69	3.31	10.69	12.69
2000 mg/kg	5	157.27	9.61	156.27	158.26	11.12	3.9	10.12	12.11
4000 mg/kg	5	143.48	20.2	142.49	144.48	10.58	4.3	6.1	17.0
DW (10 mg/kg)	5	151.25	75.3	150.03	152.47	13.4	8.3	11.2	15.6

KEY: DW = Distilled water. IU = international units. N = sample size. SD = Standard deviation. CI = confidence interval. LL= Lower limit. UL= Upper limit

Table 3. Showing descriptive statistics in body weight and serum glucose levels after two weeks

Variables	N	Mean	SD	95% Confidence Interval	
				LL	UL
Blood glucose (mMol/dl)	30	10.97	4.561	10.54	11.40
Body weight (g)	30	148.13	28.38	147.70	148.56

KEY: SD = Standard deviation; N = number. LL/UL = Lower/Upper limit. Pearson correlation = -0.385 with a P = 0.030. Spearman's correlation coefficient = -.412. P = 0.019

been seen in a previous study (primary communication) and the complete survival of the test animals would be due to its potent antioxidant properties [24]. The phytochemical analysis of the *Musa paradisiaca* extract showed the presence of alkaloids, glycosides, steroids, saponins, tannins, flavonoids and terpenoids. It also contains DPPH (1, 1-diphenyl-2-picrylhydrazyl) a free radical scavenger [25].

Bearing in mind that T1DM has no ideal therapeutic option, the study has been able to provide a scientific argument for the use of banana peels in the management of DM in rural communities that consume plenty of bananas [10]. It is worthy of note that the treatment groups were highly effective at higher dosages (Treatment > 200 mg/kg; $P \leq 0.05$). This condition would subsequently play a crucial role in reducing the estimated global and national burden of DM over time [7]. Hence would inevitably reduce the healthcare burden, thus saving the central government much money in procuring drugs and above all, reducing the rate of development of insulin resistance in the DM population [26].

The 4000 mg/kg had a more potent hypoglycemic effect than the 2000 mg/kg which may be attributable to the relatively higher concentration of the phytochemicals in the extract. Impressively, no toxicological effects were associated with the high dosage (primary observations), which we found to be interesting. Unlike other ethno medicinal plants being used on the continent to manage DM [27], it appears that *Musa paradisiaca* is a leading candidate with minimal side effects [13]. However, histopathological studies will be necessary for a better understanding of the implications of cooking the unripe banana with the peels.

In all the experimental groups, there were no significant differences ($P > 0.05$) in weight after fourteen days as when compared to the mean body weights on day 0 thus showing an efficient modulation of weight change with banana peels.

This finding supports the findings of previous studies that *Musa paradisiaca* has strong antidiabetic properties [25] and efficient in the management of body weight in DM. The preceding shows that *Musa paradisiaca* has additional pharmaceutical advantages over insulin and other medications, with weight gain implications [19]. Moreover, the study showed that 2000 mg/kg and 4000 mg/kg managed to maintain the body weights effectively, thus showing that the banana peel is a potential candidate for further drug development studies. More so, banana is associated with a high rate of cholesterol degradation [28], which is vital for DM patients in the light of cardio-vascular complications associated with DM.

Phytochemicals are found naturally in plants, and they are responsible for colour and organoleptic properties, protection. Previous reports indicated that phytoconstituents in fruits and vegetables may reduce the risk of many diseases, possibly due to dietary fibres, polyphenols, antioxidants, as well as anti-inflammatory and hepatoprotective effects [10]. Phytochemical screening of the aqueous extract of the banana peel revealed the presence of some secondary plant metabolites. These include saponins, tannins, phenols, flavonoids, cardiac glycoside, alkaloids, steroids and terpenoids, as shown in Table 1. These phytochemicals have shown hypoglycemic effects in previous studies [26].

5. CONCLUSION

In the present study, aqueous extracts of green banana peel efficiently regulated diabetic hyperglycemia and weight gain in male Wister rats at concentrations of 2000 and 4000 mg/kg body weight. It therefore, suggests an urgent change in the method of preparing *Matooke* (banana) in Uganda, which involves total peeling before boiling. In the light of the foregoing, it is recommended that *Matooke* should be boiled with the peels. This method would allow the natural phytochemicals seep into the fleshy fruit and thus deliver its therapeutic value on

consumption. Furthermore, contains phytochemicals with potentials of increasing we suggest that the individual phytochemicals be isolated and quantified in order to enable possible identification and development of drug candidates.

ETHICAL APPROVAL

Approval for animal studies were obtained from the School of Health Sciences Research Ethics Committee of Kampala International University.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO. Global report on diabetes. 2016;25. Available:www.who.int/iris/bitstream/10665/204871/1/9789241565257_eng.pdf
2. Petersen M, Yiannoutsos CT, Justice A, Egger M. Observational research on NCDs in HIV-positive populations: Conceptual and methodological considerations. *J Acquir Immune Defic. Syndr.* 2014;67:S8-16. DOI: 10.1097/QAI.0000000000000253
3. Mudie K, Jin M M, Tan K L, Addo J, Dos-Santos-Silva I, Quint J, Smeeth L, Cook S, Nitsch D, Natamba B, Gomez-Olive FX, Ako A, Perel P. Non-communicable diseases in sub-Saharan Africa: A scoping review of large cohort studies. *Journal of Global Health.* 2019;9(2):020409. Available:https://doi.org/10.7189/jogh.09.020409
4. Kataria U, Schilar D, Kumar H, Cchikara P. Cutaneous manifestations of diabetes mellitus in controlled and uncontrolled state. *International Archives of Integrated Medicine.* 2015;2(2):90-93.
5. Davison LJ. Diabetes mellitus and pancreatitis - cause or effect? *Journal of Small Animal Practice;* 2015. Available:http://doi.org/10.1111/jsap.12295
6. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2011;34(Suppl 1):62-9. Available:http://doi.org/10.2337/dc11-S062
7. Gulabani M, John M, Iseac R. *Indian Journal of Community Medicine.* 2008; 33(3):204-206. Available:https://dx.doi.org/10.4103%2F0970-0218.42068
8. Kibiringe D, Mwebaze R. Vitamin B12 deficiency among patients with Diabetes mellitus: Is routine screening and supplementation justified? *Journal of Diabetes and Metabolic Disorders.* 2013; 12:17. Available:https://dx.doi.org/10.1186%2F2251-6581-12-17
9. Mondo CK, Otim MA, Akol G, Musoke R, Orem J. The prevalence and distribution of non-communicable diseases and their risk factors in Kasese district, Uganda. *Cardiovascular Journal of Africa.* 2013; 24(3):52-7. Available:http://doi.org/10.5830/CVJA-2012-081
10. Ssenyange CW, Namulindwa A, Oyik B, Ssebulibwa J. Plants used to manage type II diabetes mellitus in selected districts of central Uganda, *African Health Science.* 2015;15(2):496-502. Available:https://dx.doi.org/10.4314%2Fahs.v15i2.24
11. Adewoye EO, Taiwo VO, Olayioye FA. Anti-oxidant and anti-hyperglycemic activities of *Musa sapientum* root extracts in alloxan-induced diabetic rats *African Journal of Medicine and Medical Sciences.* 2009;38:109-117.
12. Hussain A, Khan MN, Iqbal Z, Sajid MS, Khan MK. Anthelmintic activity of *Trianthema portulacastrum* L. and *Musa paradisiaca* L. against gastrointestinal nematodes of sheep. *Veterinary Parasitology.* 2011;179(1-3):92-99. Available:http://doi.org/10.1016/j.vetpar.2011.02.022
13. Nirmala M, Girija K, Lakshman K, Divya T. Hepatoprotective activity of *Musa paradisiaca* on experimental animal models. *Asian Pacific Journal of Tropical Biomedicine.* 2012;2(1):11-15. Available:http://doi.org/10.1016/S2221-1691(11)60181-0
14. Jawla S, Kumar Y, Khan MSY. Antimicrobial and antihyperglycemic activities of *Musa paradisiaca* flowers. *Asian Pacific Journal of Tropical Biomedicine.* 2012;2(SUPPL. 2). Available:http://doi.org/10.1016/S2221-1691(12)60336-0
15. Kappe VD, Cazarolli LH, Pereira DF, Postal BG, Madoglio FA, Buss Z da S, Silva FRMB. Beneficial effects of banana leaves (*Musa paradisiaca*) on glucose

- homeostasis: Multiple sites of action. Brazilian Journal of Pharmacognosy. 2013; 23(4):706–715.
Available:<http://doi.org/10.1590/S0102-695X2013005000062>
16. Alkarkhi AFM, Ramli S, Bin Yong YS, Easa AM. Comparing physicochemical properties of banana pulp and peel flours prepared from green and ripe fruits. Food Chemistry. 2011;129(2):312–318.
Available:<http://doi.org/10.1016/j.foodchem.2011.04.060>
17. Harborne JB. Phytochemical methods. A guide to modern techniques of plant analysis; 3rd Edition, London, Chapman and Hall. 1998;279.
18. Campos KE, Diniz YS, Cataneo AC, Faine LA, Alves MJ, Novelli EL. Hypoglycemic and antioxidant effects of onion, *Allium cepa*: Dietary onion addition, antioxidant effects on diabetic rats. International Journal of Food Sciences and Nutrition. 2003;54(3):241-246.
Available:<https://dx.doi.org/10.1080/09637480120092062>
19. Mitri J, Hamdy O. Diabetes medications and body weight. Expert Opinion on Drug Safety. 2009;8(5):573–584.
Available:<http://doi.org/10.1517/14740330903081725>
20. Liu L, Yin X, Morrissey S. Global variability in diabetes mellitus and its association with body weight and primary healthcare support in 49 low- and middle-income developing countries. Diabetic Medicine. 2012;29(8):995–1002.
Available:<http://doi.org/10.1111/j.1464-5491.2011.03549.x>
21. Tonstad S, Butler T, Yan R, Fraser GE. Type of vegetarian diet, body weight, and prevalence of type 2 diabetes. Diabetes Care. 2009;32(5):791–796.
Available:<http://doi.org/10.2337/dc08-1886>
22. Koh-Banerjee P, Franz MV, Sampson L, Liu SM, Jacobs DR, Spiegelman D, Willett W, Rimm E. Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y weight gain among men. Am. J. Clin. Nutr. 2004;80:1237–1245.
23. Anderson JW, Kendall CWC, Jenkins DJA. Importance of weight management in type 2 diabetes: Review with meta-analysis of clinical studies. College of Nutrition. 2003; 22(5):331–339.
Available:<http://doi.org/10.1080/07315724.2003.10719316>
24. Mahmood A, Ngah N, Omar MN. Phytochemicals constituent and antioxidant activities in *Musa paradisiaca* flower. European Journal of Scientific Research. 2011;66(2):311–318.
25. Hince IN, Hartwell HJ, Feng Y, Theve EJ, Hall GA, Hashway S, Conolly J, Fecteau M, Fox JG, Rogers AB. Insulin resistance and metabolic hepatocarcinogenesis with parent-of-origin effect in AxB mice. American Journal of Pathology. 2011; 179(6):2855-65.
DOI: 10.1016/j.ajpath.2011.08.014
26. Ajagun-Ogunleye OM, Tirwomwe M, Mitaki RN, Ejekwumadu JN, Kasozi IK, Pantoglou J, Mitaki NB. Hypoglycemic and high dosage effects of *Bidens pilosa* in type-1 diabetes mellitus. Journal of Diabetes Mellitus. 2015;5:146–154.
27. Iweala EE, Obichi IC, Omotosho OE. Biochemical and histological responses of hepatotoxic rats fed *Musa paradisiaca* L. - Supplemented diet. International Journal of Pharmacology. 2011;4:471-477.
28. Lamba SS, Buch KY, Lewis J. Phytochemicals as potential hypoglycemic agents. Studies in Natural Products Chemistry. 2000;21:457-495.

© 2020 Iwueke et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/57725>