

First record of baobab (*Adansonia digitata* L.) in Uganda

Jens Gebauer · Cory W. Whitney ·
John R. S. Tabuti

Received: 7 September 2015 / Accepted: 2 March 2016 / Published online: 17 March 2016
© Springer Science+Business Media Dordrecht 2016

Abstract The baobab (*Adansonia digitata* L.) is a remarkable key tree species with different uses in many African countries. International interest in the species has intensified in recent years. Despite the wide distribution of the baobab in many African regions, the tree has been reported to be absent in Uganda. In 2015 and 2016, research trips in the Central, Western, Eastern and Northern Regions of Uganda were conducted to screen the cultivated and natural flora for baobabs. As a result, four vigorous baobab trees aged 9 and 22 years were identified in two gardens in the Iganga and Soroti Districts in the Eastern Region of Uganda. The origins of the planting materials were Nyala in Sudan and Mombasa in Kenya. To our knowledge, these are the first scientific records of baobab in Uganda and some research questions and key propositions are formulated based on this discovery.

Keywords *Adansonia digitata* · Baobab · East Africa · Fruit tree · Malvaceae · Uganda

Introduction

The baobab (*Adansonia digitata* L., Malvaceae, subfamily Bombacoideae, Baum et al. 2004) also known as Africa's wooden elephant (Gebauer et al. 2016) is a remarkable key tree species in many African countries (e.g. Jama et al. 2008; Buchmann et al. 2010, Venter and Witkowski 2013). This impressive tree (for botanical description see Gebauer et al. 2002) is of economic and cultural importance. It is used for human food, water storage, fodder and medicine among many other purposes (Wickens and Lowe 2008 and references therein). In West Africa, both baobab fruits and leaves contribute to human diets in a major way, while in East and Southern Africa the fruits are the most important plant part used (NRC 2006). In recent years, international interest in the species has intensified following the acceptance of baobab fruit pulp as a food ingredient by the European Union (EC 2008) and the US Food and Drug Administration (FDA 2009). A recent study by Gebauer et al. (2014) revealed that more than 300 baobab products or products with baobab as an ingredient are available in Europe.

In Africa, products of the multipurpose tree are mainly collected from the wild and we know of no commercial baobab plantations. Due to the high local,

J. Gebauer (✉) · C. W. Whitney
Faculty of Life Sciences, Rhine-Waal University of Applied Sciences, Marie-Curie-Str. 1, 47533 Kleve, Germany
e-mail: jens.gebauer@hochschule-rhein-waal.de

C. W. Whitney
Faculty of Organic Agricultural Sciences, University of Kassel, Steinstr. 19, 37213 Witzenhausen, Germany

J. R. S. Tabuti
College of Agricultural and Environmental Sciences, Makerere University, P.O. Box 7062, Kampala, Uganda

national and international demand of baobab products, trees are intensively harvested, which often leads to overexploitation of the natural stands (Dhillon and Gustad 2004). This situation calls for the development and implementation of sustainable utilization methods and the cultivation of baobabs within and outside of its natural range. In some East and West African countries such as Niger and Sudan, the baobab can be found in urban and rural gardens (Bernholt et al. 2009; Gebauer 2005; Goenster et al. 2011; Wiehle et al. 2014). In Kenya and Tanzania baobab trees are often left standing and are protected on farms (Beentje 1989, 1994; Ruffo et al. 2002).

Despite the wide distribution and importance of the baobab in many African regions, the tree is widely known to be absent in Uganda (e.g. Katende et al. 1995; Bosch et al. 2004; Wickens and Lowe 2008; Orwa et al. 2009; Nussinovitch 2010; Dharani 2011; Kalema and Beentje 2012; Lim 2012). Ecological niche modelling results of Cuní Sanchez et al. (2010)

indicated that the presence of baobab trees is mainly related to annual precipitation and temperature seasonality. Jensen et al. (2011) stated that the ability of the species to compete with other plants is relatively weak in the humid sites. In Uganda the baobab is not only regarded as absent in natural habitats, it is also not known to be cultivated on farm-land (Eilu et al. 2003, 2007; Agea et al. 2007; Okullo and Waithum 2007; Tabuti et al. 2010; Tabuti 2012; Nyamukuru et al. 2015) or in private/homegardens (Oduol and Aluma 1990; Okullo et al. 2003; Whitney and Gebauer 2014).

Materials and methods

In July and August of 2015 and January of 2016, research trips to the Central, Western, Eastern and Northern Regions of Uganda were conducted within the collaborative research project “Reduction of Post-Harvest Losses and Value Addition in East African




Fig. 1 Map of Uganda indicating the study locations and the places of the baobab records

Food Value Chains” (RELOAD). Cultivated and natural flora was screened and local people interviewed about the occurrence of baobabs. The expeditions comprised the following main study locations: Atiak, Buhoma, Busembatia, Entebbe, Fort Portal, Gulu, Ishasha, Kamion, Kampala, Kamudini, Kasese, Kigumba, Kitgum, Kotido, Lira, Lutoto, Masaka, Mbale, Mbarara, Mubende, Nadunget, Nakosongola, Ntungamo and Soroti (Fig. 1). In total the trips covered a distance of 4975 km with a range in elevation between 809 m a.s.l. and 1680 m a.s.l. To access cultivated plant species, homegardens and fields were visited and farmers interviewed. In the surroundings of the locations and along roads, natural vegetation was screened in detail on the spot or from outcrops using Swarovski CL 8 × 30 and Leica Ultravid 10 × 25 BR binoculars. Baobab tree height and stem circumference at 130 cm was measured using a 30 m hand-operated tape. For better comparison with other studies, stem diameter at breast height

(DBH) was determined arithmetically, assuming a circular stem shape. The location of each tree was recorded using a Global Positioning System (GPS) receiver (Garmin eTrex 30).

Results and discussion

As a result of our expeditions, four vigorous baobab trees were identified in the Eastern Region of Uganda.

Two baobab trees (Fig. 2) were located to the north-east of Soroti town (Fig. 1) in the Soroti District in 2015. The trees (locations: 01°45.048'N, 33°37.010'E, 01°45.055'N, 33°37.015'E, 1070 m a.s.l.) with heights of 4.36 and 4.79 m and DBHs of 52 and 40 cm were both found at the roadside edge of a unfenced grass courtyard along Moroto Road. The garden owner told us that he obtained two seedlings from a tree nursery in Nyala in South Darfur, Sudan and planted them as ornamentals in 2007. The plants developed well,




Fig. 2 Baobab trees (a, b) and leaves (c, d) in a private garden north-east of Soroti town in the Soroti District, Eastern Region of Uganda

showing high vitality and fast growth. The garden owner had worked from 2005 to 2008 in development projects in Darfur and was impressed by the beauty of baobabs in this region. He was curious to explore the development and performance of the baobabs in Soroti's environmental conditions, with annual rainfall of between 1100 and 1200 mm. However, the trees had to be trimmed some years ago to allow construction of power lines close to the baobabs. In general, the peculiar tree shape has attracted many people's attention and is the main reason why African baobabs are planted as ornamentals throughout the tropics and subtropics (Sidibé and Williams 2002). Due to its unique eye-catching profile it is also cultivated as a bonsai in many countries (Gebauer et al. 2014).

Two more baobab trees standing side by side (Fig. 3) were recorded south-west of Busembatia town (Fig. 1) in the Iganga District in 2016. The trees (location: 00°46.148'N, 33°37.159'E, 1089 m a.s.l.)

with heights of 9.90 and 9.55 m and DBHs of 79 and 85 cm were found in a fenced homegarden along Tirinyi Road. The garden owner told us that he extracted the seeds from baobab fruits which were sent by his sister from Mombasa, Kenya and planted them in 1994. Two seeds germinated and the seedlings developed well, showing high vitality and fast growth. According to the owner, the main reason for raising the baobabs in his garden was to harvest fruits. He was impressed by the delicious taste of the baobab fruits and was hopeful that those he planted would bear fruits. However, so far the 22 year old baobabs have not borne any fruits. There is little information available on the age at which baobabs first flower. Wickens (1982) and Assogbadjo and Loo (2011) noted that this can take up to 23 years. According to Watson (2007) it can take 30 years for a baobab to produce the first flowers. Therefore, efforts are made to accelerate fruit precocity through horticultural techniques such




Fig. 3 Baobab trees (a) and leaves (b) in a homegarden south-west of Busembatia town in the Iganga District, Eastern Region of Uganda



Fig. 4 Baobab tree (a) and leaves (b) in Entebbe Botanical Garden in the Wakiso District, Central Region of Uganda

as grafting (Kalinganire et al. 2007, Anjarwalla et al. 2015).

To our knowledge these are the first scientific records of baobab in Uganda. An extensive search of literature yielded only one internet source (Geoview.info 2007) displaying a photo of a large mature baobab, claiming that this picture was taken in Entebbe, Central Region of Uganda. We searched the geolocation and the town of Entebbe extensively and did not find any sign of a baobab. People living in the area also stated that this photo was not of a tree they had ever seen.

The extensive collection of Entebbe Botanical Garden in the Central Region of Uganda has one stunted baobab (location: 00°03.785'N, 32°28.643'E, 1158 m a.s.l.). This tree (Fig. 4) is claimed to be more than 100 years old (pers. comm. with Entebbe Botanical Garden guide 2016). Yet it has a height of just 4.20 m and a DBH of only 13 cm. The Botanical Garden has no record of flowering for the tree. Reasons for the weak performance of the tree can be seen in the

tropical wet climate with no significant dry or cold season with a year-round rainfall of around 1600 mm. According to von Maydell (1990) the species grows best in regions with an annual rainfall of between 250 and 1000 mm. In contrast to Entebbe, the much dryer climatic conditions in Busembatia and Soroti with prolonged drought followed by heavy rain probably led to the vigorous and fast growth of the recorded baobabs.

Research perspectives

Based on the findings of this preliminary study the following research questions arise:

- How will these baobab trees perform in the future?
- When will they start to flower and fruit?
- What fruit quality can be expected and will local people accept it as food?
- Will this tree species become a common plant of homegardens in the Eastern Region of Uganda?

Are there more unknown baobabs in homegardens, on farm-land and/or in the wild of Uganda?

In conclusion it would be important to monitor the performance of the four baobab trees of the two geographical origins in the coming decades and to establish provenance trials including accessions from Western and Southern Africa for evaluation in different regions of Uganda. Such trials can show remarkable differences in survival and performance between provenances (Ræbild et al. 2011; Korbo et al. 2012; Munthali et al. 2012). According to Cuní Sanchez et al. (2010, 2011) it is worthwhile to identify potential cultivation areas outside of the species current distribution range to set up ex situ conservation sites and to support sustainable production of baobab products in Africa. This would be important to satisfy the growing African and international demand for products such as baobab fruit pulp, which is a key raw material for the wide range of baobab products or products that contain baobab as an ingredient (Gebauer et al. 2014). The screening of rural and urban markets for baobab products in different regions of Uganda could determine whether baobab products are available e.g. from neighbouring countries, such as South Sudan, Kenya and/or Tanzania, where the species grows naturally and baobab products are common (El Amin 1990; Murray et al. 2001; Ruffo et al. 2002; Robinson 2004; Mwema et al. 2012; McMullin and Kehlenbeck 2015). This would offer insights regarding the cultural acceptance of baobab products for food and nutritional security and other domestic purposes. Furthermore, in-depth botanical studies of wild and cultivated flora with a particular emphasis on baobab identification and evaluation in the dry northern part of the country may produce further evidence of the local presence of the baobab, Africa's wooden elephant in Uganda.

Acknowledgments This study was part of the subproject 4, work package 2 (031A247B) within the collaborative research project “Reduction of Post-Harvest Losses and Value Addition in East African Food Value Chains” (RELOAD) of the initiative “GlobE—Research for the Global Food Supply” of the German Federal Ministry of Education and Research (BMBF). We are also grateful to Antonia Nyamukuru, PhD candidate at Makerere University in Kampala, Uganda and Sebastian Walaita Javan affiliated with Delight Uganda Ltd. in Kampala, Uganda for collecting information on promising study locations. To Richard Bawonga we are thankful for driving us safely all the 4975 km through Uganda. Furthermore, we also acknowledge the support of Dr Eike Luedeling affiliated with the World Agroforestry Centre (ICRAF) in Nairobi, Kenya and the Center for

Development Research (ZEF) in Bonn, Germany for creating the Uganda map (Fig. 1).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Agea JG, Obua J, Kaboggoza JR, Waiswa D (2007) Diversity of indigenous fruit trees in the traditional cotton-millet farming system: the case of Adwari Subcounty, Lira district, Uganda. *Afr J Ecol* 45:39–43
- Anjarwalla P, Ofori D, Owino A, Matuku D, Kehlenbeck K (2015) Testing different vegetative propagation methods for baobab (*Adansonia digitata* L.) to assist its domestication and promote cultivation. XIV World Forestry Congress in Durban, South Africa, pp 1–8
- Assogbadjo AE, Loo J (2011) *Adansonia digitata*, African baobab. Conservation and sustainable use of genetic resources of priority food tree species in sub-Saharan Africa. Bioversity International, Italy
- Baum DA, Smith SD, Yen A, Alverson WS, Nyffeler R, Whitlock BA, Oldham RL (2004) Phylogenetic relationships of Malvaceae (Bombacoideae and Malvoideae; Malvaceae *sensu lato*) as inferred from plastid DNA sequences. *Am J Bot* 91:1863–1871
- Beentje HJ (1989) Flora of tropical East Africa: Bombacaceae. Balkema, The Netherlands
- Beentje HJ (1994) Kenya trees, shrubs and lianas. National Museums of Kenya, Kenya
- Bernholt B, Kehlenbeck K, Gebauer J, Buerkert A (2009) Plant species richness and diversity in urban and peri-urban gardens of Niamey, Niger. *Agrofor Syst* 77:159–179
- Bosch CH, Sié K, Asafa BA (2004) *Adansonia digitata* L. In: Grubben GJH, Denton OA (eds) Plant resources of tropical Africa 2. Vegetables. PROTA Foundation. Backhuys Publishers, Netherlands
- Buchmann C, Prehlsler S, Hartl A, Vogl CR (2010) The importance of baobab (*Adansonia digitata* L.) in rural West African subsistence—suggestion of a cautionary approach to international market export of baobab fruits. *Ecol Food Nutr* 49:145–172
- Cuní Sanchez A, Osborne PE, Haq N (2010) Identifying the global potential for baobab tree cultivation using ecological niche modelling. *Agrofor Syst* 80:191–201
- Cuní Sanchez A, Osborne PE, Haq N (2011) Climate change and the African baobab (*Adansonia digitata* L.): the need for better conservation strategies. *Afr J Ecol* 49:234–245
- Dharani N (2011) Field guide to common trees and shrubs of East Africa. Struik Nature, South Africa
- Dhillion SS, Gustad G (2004) Local management practices influence the viability of the baobab (*Adansonia digitata* Linn.) in different land use types, Cinzana, Mali. *Agric Ecosyst Environ* 101:85–103
- EC (2008) Commission decision: authorizing the placing on the market of baobab dried fruit pulp as a novel food ingredient

- under regulation (EC) No 258/97 of the European Parliament and of the Council. Off J EU L 183(38):1–2
- Eilu G, Obua J, Tumuhaiwe JK, Nkwine C (2003) Traditional farming and plant species diversity in agricultural landscapes of south-western Uganda. *Agric Ecosyst Environ* 99:125–134
- Eilu G, Obua J, Tumuhaiwe JK, Nkwine C (2007) Conservation of indigenous plants outside protected areas in Tororo District, eastern Uganda. *Afr J Ecol* 45:73–78
- El Amin HM (1990) Trees and shrubs of the Sudan. Ithaca Press, UK
- FDA (2009) Agency response letter GRAS Notice No. GRN 000273, 25/July/2009. US Food and Drug Administration. <http://www.fda.gov/food/ingredientspackaginglabeling/gras/noticelabeling/ucm174945.htm>. Accessed 27 Dec 2015
- Gebauer J (2005) Plant species diversity of home gardens in El Obeid, Central Sudan. *J Agric Rural Dev Trop Subtrop* 106:97–103
- Gebauer J, El-Siddig K, Ebert G (2002) Baobab (*Adansonia digitata* L.): a review on a multipurpose tree with promising future in the Sudan. *Eur J Hort Sci* 67:155–160
- Gebauer J, Assem A, Busch E, Hartmann S, Möckel D, Krebs F, Ziegler T, Wichern F, Wiehle M, Kehlenbeck K (2014) Der Baobab (*Adansonia digitata* L.): Wildobst aus Afrika für Deutschland und Europa?! [The baobab (*Adansonia digitata* L.): wild African fruit for Germany and Europe?!]. *Erwerbs-Obstbau* 56:9–24
- Gebauer J, Adam YO, Cuní Sanchez A, Darr D, Eltahir MES, Fadl KEM, Fernsebner G, Frei M, Habte T-Y, Hammer K, Hunsche M, Johnson H, Kordofani M, Krawinkel M, Kugler F, Luedeling E, Mahmoud TE, Maina A, Mithöfer D, Munthali CRY, Noga F, North R, Owino WO, Prinz K, Rimberia FK, Saied A, Schüring M, Sennhenn A, Späth MA, Taha MEN, Triebel A, Wichern F, Wiehle M, Wrage-Mönnig N, Kehlenbeck K (2016) Africa's wooden elephant: the baobab tree (*Adansonia digitata* L.) in Sudan and Kenya: a review. *Genet Resour Crop Evol*. doi:10.1007/s10722-015-0360-1
- Geoview.info (2007) http://ug.geoview.info/baobab_tree,6250993p. Accessed 11 Aug 2015
- Goenster S, Wiehle M, Kehlenbeck K, Jamnadass R, Gebauer J, Buerkert A (2011) Indigenous fruit trees in homegardens of the Nuba Mountains, Central Sudan: tree diversity and potential for improving nutrition and income of rural communities. *Acta Hort* 911:355–364
- Jama BA, Mohamed AM, Mulatya J, Njui AN (2008) Comparing the “big five”: a framework for the sustainable management of indigenous fruit trees in the drylands of east and central Africa. *Ecol Indic* 8:170–179
- Jensen JS, Bayala J, Sanou H, Korbo A, Ræbild A, Kambou S, Tougiani A, Bouda H-N, Larsen SA, Parkouda C (2011) A research approach supporting domestication of baobab (*Adansonia digitata* L.) in West Africa. *New For* 41:317–335
- Kalema J, Beentje H (2012) Conservation checklist of the trees of Uganda. Kew Publishing Royal Botanic Gardens, UK
- Kalinganire A, Weber J, Uwamariya A, Kone B (2007) Improving rural livelihoods through domestication of indigenous fruit trees in the parklands of the Sahel. In: Akinnifesi F, Leakey R, Ajayi O, Sileshi G, Tchoundjeu Z, Matakala P, Kwesiga F (eds) *Indigenous fruit trees in the tropics: domestication, utilization and commercialization*. CAB International Publishing, USA
- Katende AB, Birnie A, Tengnas BO (1995) Useful trees and shrubs for Uganda: identification, propagation, and management for agricultural and pastoral communities. Regional Soil Consultants Unit, Kenya
- Korbo A, Sanou H, Ræbild A, Jensen JS, Hansen JK, Kjær ED (2012) Comparison of East and West African populations of baobab (*Adansonia digitata* L.). *Agrofor Syst* 85:505–518
- Lim TK (2012) Edible medicinal and non-medicinal plants: volume 1, fruits. Springer, Netherlands
- McMullin S, Kehlenbeck K (2015) Untapped potential of baobab. *Miti* 26:16–17
- Munthali CRY, Chirwa PW, Akinnifesi FK (2012) Genetic variation among and within provenances of *Adansonia digitata* L. (baobab) in seed germination and seedling growth from selected natural populations in Malawi. *Agrofor Syst* 86:419–431
- Murray SS, Schoeninger MJ, Bunn HT, Pickering TR, Marlett JA (2001) Nutritional composition of some wild plant foods and honey used by Hadza foragers of Tanzania. *J Food Compos Anal* 14:3–13
- Mwema CM, Mutai BK, Lagat JK, Kibet LK, Maina MC (2012) Contribution of selected indigenous fruits on household income and food security in Mwingi, Kenya. *Curr Res J Soc Sci* 4:425–430
- NRC (2006) Lost crops of Africa. National Research Council, vol II. The National Academies Press, USA
- Nussinovitch A (2010) Plant gum exudates of the world: sources, distribution, properties, and application. CRC Press, USA
- Nyamukuru A, Nabatanzi A, Mpiira S, Tabuti JRS (2015) Locally preferred woody species and their management in Kiruhura and Arua districts, Uganda. *Ethnobot Res App* 14. doi:10.17348/era.14.0.049-061. Accessed 06 Jan 2016
- Oduol PA, Aluma JRW (1990) The banana (*Musa* spp.)—*Coffea robusta*: traditional agroforestry system of Uganda. *Agrofor Syst* 11:213–226
- Okullo JBL, Waithum G (2007) Diversity and conservation of on-farm woody plants by field types in Paromo Subcounty, Nebbi District, north-western Uganda. *Afr J Ecol* 45:59–66
- Okullo JBL, Obua J, Kaboggoza JRS, Aluma JRW (2003) Traditional agroforestry systems, tree uses and management in Northern Uganda. *UJAS* 8:5–10
- Orwa C, Mutua A, Kindt R, Jamnadass R, Simons A (2009) Agroforestree Database: a tree reference and selection guide. Version 4.0. <http://www.worldagroforestry.org/resources/databases/agroforestree>. Accessed 27 Dec 2015
- Ræbild A, Larsen AS, Jensen JS, Ouedraogo M, De Groote S, van Damme P, Bayala J, Diallo BO, Sanou H, Kalinganire A, Kjær ED (2011) Advances in domestication of indigenous fruit trees in the West African Sahel. *New For* 41:297–315
- Robinson J (2004) After the conflict: plant genetic resources of southern Sudan. *Plant Genet Resour* 2:85–97
- Ruffo CK, Birnie A, Tengnäs B (2002) Edible wild plants of Tanzania. Regional Land Management Unit (RELMA). English Press, Kenya
- Sidibé M, Williams JT (2002) Baobab *Adansonia digitata*. International Centre for Under-utilised Crops, UK

- Tabuti JRS (2012) Important woody plant species, their management and conservation status in Balawoli Sub-county, Uganda. *Ethnobot Res Appl* 10:269–286
- Tabuti JRS, Muwanika VB, Arinaitwe MZ, Ticktin T (2010) Conservation of priority woody species on farmlands: a case study from Nawaikoke Sub-county, Uganda. *Appl Geogr* 31:456–462
- Venter SM, Witkowski ETF (2013) Fruits of our labour: contribution of commercial baobab (*Adansonia digitata* L.) fruit harvesting to the livelihoods of marginalized people in northern Venda, South Africa. *Agrofor Syst* 187:159–172
- Von Maydell H-J (1990) Trees and shrubs of the Sahel: their characteristics and uses. Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ) GmbH. Verlag Josef Margraf, Germany
- Watson R (2007) The African baobab. Struik, South Africa
- Whitney CW, Gebauer J (2014) Homegardens in Uganda: diversity and potential. In: Rahmann G, Aksoy U (eds) Building organic bridges. Proceedings of the 4th ISOFAR scientific conference at the organic world congress in Istanbul, Turkey, pp 1115–1118
- Wickens GE (1982) The baobab—Africa’s upside-down tree. *Kew Bull* 37:173–209
- Wickens GE, Lowe P (2008) The baobabs: pachycauls of Africa. Springer, Madagascar
- Wiehle M, Prinz K, Kehlenbeck K, Goenster S, Mohamed SA, Finkeldey R, Buerkert A, Gebauer J (2014) The African baobab (*Adansonia digitata* L.)—genetic resources in neglected populations of the Nuba Mountains, Sudan. *Am J Bot* 101:1489–1507