

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325787584>

Initial Efficiency of Commonly Used Practices to Control Soil, Runoff and Nutrient Losses from Maize and Banana Based Systems in the Lake Kivu Basin

Article in *Journal of Experimental Agriculture International* · June 2018

DOI: 10.9734/JEAI/2018/23636

CITATIONS

2

READS

40

5 authors, including:



Majaliwa J. G. Mwanjalolo

RUFORUM

91 PUBLICATIONS 1,071 CITATIONS

[SEE PROFILE](#)



Moses Tenywa

Agricultural innovation systems brokerage association AGINSBA

117 PUBLICATIONS 2,732 CITATIONS

[SEE PROFILE](#)




Adipala Ekwamu

RESEARCH AND EDUCATION AGENCY

196 PUBLICATIONS 2,882 CITATIONS

[SEE PROFILE](#)



Initial Efficiency of Commonly Used Practices to Control Soil, Runoff and Nutrient Losses from Maize and Banana Based Systems in the Lake Kivu Basin

M. W. Adidja^{1*}, J. G. M. Majaliwa¹, M. M. Tenywa¹, S. Bashwira² and E. Adipala³

¹*College of Agricultural and Environmental Sciences, Makerere University, P.O.Box 7062, Kampala, Uganda.*

²*Université Catholique de Bukavu, P.O.Box 285 Bukavu, South Kivu, D.R. Congo.*

³*RUFORUM, Plot 151, Garden Hill Makerere University, Main Campus Kampala, P.O.Box 7062, Kampala, Uganda.*

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2018/23636

Editor(s):

(1) Anita Biesiada, Professor, Department of Horticulture, Wrocław University of Environmental and Life Sciences, Poland.

Reviewers:

(1) Ernest Frimpong Asamoah, Beijing Normal University, China.

(2) Artemi Cerda, Universitat de València, Spain.

Complete Peer review History: <http://www.sciencedomain.org/review-history/25118>

Original Research Article

Received 11th December 2015

Accepted 22nd March 2018

Published 13th June 2018

ABSTRACT

This study was carried out to determine the initial efficiency of commonly used practices to control soil erosion from Maize and Banana based cropping systems in Lwiro micro catchment in the Lake Kivu basin. Soil, runoff and nutrient losses were determined using runoff plot approach. Instrumentalised runoff plots of 2X15m were installed on maize intercropped with beans and banana gardens. Two soil erosion management practices, namely; *Tithonia* and contour bunds were tested on Maize intercrop with Beans and mulch for Banana. The experiment included a control practice for each crop. Each treatment and control was replicated four times. Runoff and soil loss were estimated for each rainfall event and aggregated on seasonal basis. Nutrient (N, P and K) losses were estimated per season. Results of the long and short rains of the first year of experimentation show that soil and runoff losses did not significantly change with practices and seasons ($P>0.05$) for both banana and maize based systems. Soil and runoff losses ranged from 15.73 to 32.93 Mg/ha,

*Corresponding author: E-mail: adidjamatabaro@gmail.com;

and from 168.14 to 322.17 m³; respectively. Nutrient losses varied with practices and seasons (P<0.05) and ranged from 54.68 to 112.34 Kg/ha, 87.7 to 409.4 Kg/ha; 24.5 to 94.22 Kg/ha for K, N and P; respectively. Soil and runoff losses ranged from 8.99 to 20.6 t/ha, and from 85 to 152 m³; respectively. Only K losses changed significantly with season (P<0.05) and ranged from 17.8 to 53.9 Kg/ha under Banana cropping system.

Keywords: Land degradation; Lake Kivu Basin; pollution loading; Bukavu; D.R. Congo.

1. INTRODUCTION

Soil loss and land degradation are considered as challenges of global dimension [1]. Soil erosion is widespread [2-5] and varies in magnitude from place to place depending on the land-use system, population pressure, community wealth, management, relief, and vulnerability of the soil to climate aggressivity [6-8]. [3] predicted about 15 Mg/ha/yr in Kondoa Eroded Area in Tanzania. [9] observed that soil loss was 30 times on road embankments in eastern Spain than on vegetated land. [10] observed that rills on roads in south eastern region of South Africa. Their dimension was significantly determined by gradient and percentage of vegetation cover of the roadcut while their widths and depths increased with the slope gradient and decrease with percentage of vegetation cover. The diversity of species in the vegetation covers also plays a role in the magnitude of soil loss [2] and increased soil erosion might lead to reduced plant diversity. Due to increased grazing intensity soil erosion is becoming serious in the Tibetan grassland [11]. In several regions across the globe areas, steep slope with low vegetation cover including those with relatively bareland due to fire experience severe erosion [12]. In Europe, soil erosion is most prone on vineyard [13,14] and on agrosilvopastoral mountainous ecosystems [5].

On the African continents, soil erosion is also alarming. In Ethiopia, many reservoirs constructed for hydropower generation, water supply have been silted due to the alarming rate of soil erosion in some parts of Ethiopia [15]. In DRC, and particularly in Kivu mountain areas, soil erosion is believed to be one of the major processes of soil degradation [6]. The latter is reported to have reached catastrophic proportions on agricultural lands in the Kivu mountainous region [6,7,16].

Subsequently, the region is threatened by famine and food insecurity, it is reported that 70% of the population is affected by food insecurity [17]. Yet, the region has a great agricultural potential. Under rain-fed conditions, DRC can feed more

than two billion people and 50% of its agricultural potential is located in the great Kivu region (Sud-Kivu, Maniema and Nord-Kivu). This situation is likely to exacerbate and reinforce the cycle of poverty-poor management- natural resources degradation [18-20]. On the other hand, soil erosion in the Kivu mountain region contribute to increased pollution loading into Lake Kivu, one of the deadly Lakes in the world. The Lake has reached critical concentration of CH₄ and CO₂ which can lead to an increased instability of the Lake which can trigger an explosion if any action is not taken to control the flow of carbon into the water bodies [21,22]. Lake Kivu is critical for the survival of millions of people around its basin and beyond [23]. It is a biodiversity hotspot area [24] and a source of protein and income for communities around it.

Although soil erosion is considered to be a major cause of land degradation in Kivu mountain region, very few studies have been conducted in the region to determine its magnitude and evaluate practices currently used for its control. Limited studies were conducted during the colonial time and in the 80s [25], and in Rwanda [26,27,28]. Acceptability, maintenance and replicability of some of these technologies has been challenging in the Kivu. Farmers' poverty and a long time span return of these technologies being the most determining factors. For many farmers adoption and continuous investment in these practices also depend on their short time performance on yield [29] which to some extent associated with their efficiency to control soil and nutrient losses, and enhance moisture conservation in the soil [30]. This study intended to determine the initial efficiency of commonly used practices to control soil erosion from Maize and Banana based cropping systems in Lwiro micro catchment in the eastern D.R. Congo side of the Lake Kivu basin.

2. METHODOLOGY

2.1 Description of the Study Site

This study was conducted in the River Lwiro micro-catchment within Lake Kivu Basin. The

River Lwiro is located on the eastern flank of Lake Kivu between latitudes 2°15' and 2°30' S and longitudes 28°45' and 28°85' E. Its headwaters are in the Kahuzi-Biega National Park mountain region; at an altitude of 2000 m. The 84 km² river basin is bordered on the east by Lake Kivu and on the west by the Kahuzi mountain forest (Fig. 1). This watershed of Lwiro river, the principal tributary of the Lake Kivu, covered 4 localities namely Irhambi/Katana, Bugorhe, Luhihi and Bushumba in the territory of Kabare, province of South-Kivu, Democratic Republic of Congo. Annual rainfall varies between 1134 mm and 1689 mm with an average of 1411 mm. The rainfall is bimodal with a dry season from June and July. The soil comprises of clay and rich volcanic soil, which is easily eroded. The geological composition is of Precambrian metamorphoses sediments (metamorphic rocks) and Preterozoic platform sediments [31]. [32]; describes metamorphic limestone and numerous travertines along Lake Kivu and Lake Edward. Carbonates for the production of cement are also found north and north-west of Lake Kivu [7].

and elongated hills, developed on the weathered lavas. In the Bukavu area, folded and faulted Precambrian strata are covered by thick Tertiary and Quaternary lava flows. The oldest series, not present at Bukavu itself, predates local rifting and is dated between 7 and 10 millions of years [34]. The middle and upper series are present at Bukavu. The middle series, of Mio-Pliocene age [35,34], is intimately related to the rift faults. The upper series started during the Pleistocene and continued to the last century. The chemical composition of these three series evolves from sub-alkaline over moderately alkaline to strongly alkaline [36]. Every lava series consists of many individual flows, separated in time [37]. The complex geometry of the present lava layers resulted from successive rifting and eruptive episodes. Weathering and erosion, as well as normal faulting, occurred between successive lava flow series, explaining the occurrence of palaeo-relief, contact metamorphism, smectite layers, probably Vertic palaeosoils, and clastic deposits having an alluvial and colluvial origin.

2.2 Design, Treatments and Replications

The micro-catchment is dominated by small scale farming, forest and built-up area. Forest cover is more located in the mountainous area and is part of the Kahuzi Biega National Park. The geology of Lake Kivu has been described by [33] and [28]. It is a rolling landscape of convex

Soil, runoff and nutrient losses were determined using runoff plot approach. All the plots were established on a slope of 15%. A total of 12 Plots measuring 2X15 m were established on a farmer's garden at "Centre de Recherche en Science Naturelles" (CRSN-Lwiro) used for




Fig. 1. Lwiro micro-catchment and the main rivers, Lake Kivu basin




Fig. 2. Annual rainfall in the Lwiro micro-catchment
(Source Merra Data)




Fig. 3. Monthly rainfall in Lwiro micro-catchment
(Source Merra data)

growing maize and beans. Eight (8) other plots were established on a banana plantation. Each of the plots was equipped with dividers and collecting tanks. Two treatments and one control were imposed on the maize intercropped with beans garden; namely *tithonia* and contour bunds. On the banana garden one treatment (mulch) and a control were imposed. The runoff transfer coefficients were estimated in the field using water and an accurate balance. Local varieties of maize, beans and banana were used for the experiment; and banana was cultivated in monoculture. Each treatment and control was replicated four times.

2.3 Measured Parameters

The measured parameters included soil loss, runoff, nutrient losses and the crop yield. Runoff and soil loss were estimated for each rainfall event. Samples of collected runoffs were measured using a graduated cylinder and total runoff for the rainfall events were estimated by multiplying the collected runoff by the deviser's transfer coefficient. The collected runoff was thoroughly mixed and a 100 ml sample was collected for sediment concentration determination in the laboratory. In the laboratory, sediment concentration was determined by

filtration. The total soil loss for the rainfall event was determined by multiplying the total runoff volume by the associated sediment concentration. Seasonal runoff was computed as the sum of the different event runoff [38]. The same was done for soil loss. Seasonal nutrient (N, P and K) losses were estimated for the short rainy season of year 2012 and the long rainy season of the year 2013. Plot composite soil samples were obtained by putting together sediment collected after each rainfall event for a given season. The composite samples were taken to the laboratory for N, P and K analysis. N, P and K were analysed using standard procedures [39]. Total N were analysed using Kjeldahl digestion method. Available P was extracted by Bray II method [40]. The exchangeable K⁺ cation was determined by a flame photometer. Total N and total available P was determined by multiplying their respective concentration with the total season soil loss.

2.4 Data Analysis

The survey questionnaires were coded and entered in PASW statistics version 18 for analysis. Frequencies, tables and graphs were obtained using descriptive statistics and cross tabulation. For the experiment, Analysis of Variance was used for mean separation at $p \leq 0.05$ in GenStat 13th edition.

3. RESULTS

3.1 Soil and Runoff Losses under Maize Based Cropping System

Soil and runoff losses for both season I (long rainy) and season II (short rainy) are presented in –Figs. 4 and 5 respectively. Soil loss and runoff losses did not significantly depend on soil erosion practices and rain seasons ($P > 0.05$). However, soil loss under contour bunds was relatively the lowest followed by *Tithonia* and control for both seasons. A similar trend to that of soil loss was observed for runoff. The long rains season tended to have relatively higher value of soil loss and runoff for all practices. Soil loss under contour bunds varied between 15.73 t/ha and 20.49 t/ha, 19.29 t/ha and 23.33 t/ha under *Tithonia* and 30.91 and averaged 32.93 t/ha/season for the control for the two seasons. Runoff under contour bunds was also the lowest followed by *Tithonia* and control for both seasons. Runoff ranged between 168.14 and 203.22 under contour bunds, 206.60 and 240.13

under *Tithonia* and 264.29 and 322.17 t/ha/season under control.

Table 2 shows nutrient concentration under different practices and across seasons. Different nutrients were affected differently by practices and seasons. Practices effects was observed only on K while seasons affected N and P concentration in the sediment lost ($P \leq 0.05$). The interaction between seasons and practices was significant for K and P concentration in sediment lost ($P \leq 0.05$). Contour bunds tended to have a relatively high concentration of N and K and lower value of P compared to other technologies for the short rain season. All technologies had less variation in concentration of nutrients during the long rain season. Nitrogen concentration in sediments was relatively higher during the first season compared to the second season for all practices. Phosphorous concentration was relatively constant under *Tithonia* and control during the two seasons. It is important to observe that P concentration increased during the second season for contour bunds. K concentration increased under *Tithonia* and control during the second season. It is also important to note that K concentration decreased under contour bunds during the season.

Table 3 shows nutrient losses under different practices and across seasons. Nutrient losses from different land-use varied with practices and seasons ($p \leq 0.05$). Practices effects was observed on K, while N loss was only affected by season ($P \leq 0.05$). K loss and P loss was also affected by season ($P < 0.05$). The interaction between seasons and practices was significant for K and P losses ($P < 0.05$). Contour bunds tended to have a relatively high amount of K and N losses and lower value of P loss compared to other technologies for the short rainy season ($P < 0.05$). Contour bunds had also relatively low amount of K loss, N loss and P loss compared to other technologies in the long rain season ($p < 0.05$). The amount of K and N losses decreased for plots with contour bunds during the long rain season while it increased for P loss. The amount of K loss and P loss increased for control and *Tithonia* during the long rain season while N loss decreased.

3.2 Soil and Runoff Losses under Maize Based Cropping System

Soil and runoff losses for both season I (long rainy) and season II (short rainy) are presented

in Figures 6 and 7 respectively. Soil loss and runoff losses did not significantly depend on land-use practices and rainy seasons ($P>0.05$). However soil loss on mulched banana was relatively the lowest compared to the control (banana unmulched) for both seasons. A similar trend to that of soil loss was observed for runoff. The long rainy season tended to have relatively higher value of soil loss and runoff for both

practices. Soil loss under mulched banana varied between 89.93 and 15.77 t/ha/season and between 13.88 and 20.61 t/ha/season under unmulched (banana). Runoff under mulched banana was also the lowest compared to the unmulched banana. Runoff varied also between 85 and 134 under mulched banana and between 118 and 152 t/ha/season under unmulched banana.




Fig. 4. Soil losses under Maize cropping system




Fig. 5. Runoff under maize cropping system

Table 1. Nutrient concentration under different practices and across seasons

Land use practices	Short rainy 2012			Long rainy 2013		
	N	P	K	N	P	K
	%					
Control	0.87	0.23	0.19*	0.36	0.28	0.34*
Contour bunds	1.94	0.16*	0.58	0.38	0.28*	0.36
<i>Tithonia</i>	0.75	0.24	0.20*	0.4	0.26	0.36*
LSD-landuse	NS	0.06	0.34	NS	NS	NS
LSD Season	0.56					
LSD Landuse*season	NS 0.06 0.23					

*: Seasonal effects at $P < 0.05$; NS = Not significant




Fig. 6. Soil losses under banana cropping system




Fig. 7. Runoff under banana cropping system

Table 2. Nutrient losses under different practices and across seasons

Land use practices	Short rainy 2012			Long rainy 2013		
	N	P	K	N	P	K
	Kg/ha/season					
Control	291.8	68.01	54.68	125.9	94.22	112.34
Contour bunds	409.4*	24.5	103.74	71.5*	55.5	72.92
<i>Tithonia</i>	131.1	45.47	35.96	87.7	59.68	83.06
LSD- landuse	NS	25.26	28.81	NS	NS	NS
LSD Season	178.3	20.45	23.32			
LSD landuse*season	311.2	NS	40.72			

* Seasonal effects at $P < 0.05$ **Table 3. Nutrient concentration under different practices and across seasons for Banana**

Land use practices	%					
	Short rainy 2012			Long rainy 2013		
	N	P	K	N	P	K
Banana unmulched	0.74	0.24	0.19	0.39	0.23	0.28
Banana mulched	0.55	0.26	0.21	0.42	0.24	0.32
LSD- landuse	0.08	NS	NS			
LSD-season	0.07	NS	0.04			
LSD landuse.season	0.11	NS	NS			

Table 4. Nutrient losses under different practices and across season for Banana

Practices	Kg/ha/season					
	Short rainy 2012			Long rainy 2013		
	N	P	K	N	P	K
Banana unmulched	107	35.2	25.7	79	48.1	53.9
Banana-mulched	46	23.6	17.8	69	37.6	51.6
LSD-landuse	NS	NS	NS			
LSD Season	NS	NS	11.3			
LSD Landuse.Season	NS	NS	NS			

Table 4 shows nutrient concentration under different practices and across seasons. N and P were not significantly affected by different practices and seasons ($P > 0.05$). Only K was affected by the season ($P < 0.05$). All the nutrients tended to have relatively the same concentration during the long rainy season for the different practices except the N concentration which decreased under control (banana unmulched) during the long rain season. Table 7 below shows nutrient losses under different practices and across seasons. Most of the nutrient losses were not affected significantly by practices and seasons ($P > 0.05$). Only K losses were significantly affected by the season ($P < 0.05$). Mulched banana plots tended to have a relatively low nutrient losses in both seasons. K and P losses increased during the long rainy season for all practices while N losses decreased during the long rainy season for the unmulched banana.

Fig. 8. shows monthly rainfall amount in the catchment during the two seasons. Peak rainfall amount were observed in October 2012 and December 2012. Monthly rainfall amount remained less than 150 mm from february to April 2013.

4. DISCUSSION OF RESULTS

The annual soil and runoff losses from both maize and banana cropping systems are severe and moderate; respectively [41], [42]. Nutrient losses are relatively lower under Banana compared to Maize. Soil and water management practices effects were not felt during the first year of experimentation for soil and runoff but was significant for nutrient losses under Maize and only for K under banana ($P < 0.05$).

Soil loss under maize and banana in the study area are comparable with results obtained in the region [43-45]. [43] obtained soil losses of 45 t/ha

on maize intercrop with beans under sub-humid conditions and on slopes of 22% in the Lake Victoria crescent. Measured average annual soil loss ranged from 86.8 t/ha/yr in Rakai; and 27.9 t/ha/yr under banana [44]. Similar results were observed by [46] where annuals lost 71 t/ha/season and 25.1 t/ha/season for banana. Runoff observed under this study was relatively very low compared to values obtained by [44] for the same land use. Runoff under maize and banana was a seventh and half of what was obtained by [44] for the same land use respectively. Nutrient losses were very high but less than values obtained by [44] in Rakai. Values of 0-2 and 0.1-13 kg ha⁻¹ season⁻¹ for P and K, respectively, were reported on arable lands of sub-Saharan Africa [47]. A study by [48] showed that mulch could reduce up to 46% of runoff, and soil losses up to 88.7% and sediment concentration up to 84.4 % through increased infiltration. They observed that infiltration rate increased on average up to 167 % with mulch. In ants infested up to 167 % with mulch. In ants infested areas, up to 167 % with mulch. In ants infested areas, application of mulch materials can attract increased ants activities which will increase the macro pores and hence water infiltration [49].

Differences between the results of this study and aforementioned ones are attributed to slope, soil type and climatic conditions. However, very small

values (1.4 t/ha/yr) of soil loss were measured by [50] in Kabale for similar slope of 10 % and quasi similar values of 30-t ha⁻¹ yr⁻¹ were recorded on slopes of 20%. Type of soil affects soil loss through infiltration and erodibility. The infiltration rate of the experimental soil was generally very rapid [51] compared to moderate and rapid rate under [44] and [52] soils. High infiltration rate contribute to reduced runoff, runoff sediment detachment [53] and nutrient loss [54], [55], [56]. Several studies demonstrate that runoff generation is driven by antecedent soil moisture [57-60], and soil management systems [61,62]. The latter generally affects soil surface structure [63,64], [53]. A strong relationship exists between soil structure, soil water retention and organic matter [65]. The experimental soils are deep and have high organic matter contents ranging from 3.68 to 5.7 %. This explains the higher values of infiltration for these soils.

Variation in runoff, soil loss and nutrient losses between the different studies is also due to aggressive climatic factors. The erosivity factor is relatively higher in the study area than in Uganda where Mulebeke and Majaliwa conducted their studies [44,52,25]. It was ranging from 170 and 196 J m² in Uganda while in DR Congo it ranged between 220 and 300 J m². This explain the need of soil and water conservation practices to control soil erosion which could be induced by such rainfall aggressivity [66,14,67].




Fig. 8. Rainfall data during the study period

It is worthwhile to note that maize and beans yields were generally high despite declining in the second season. In Ethiopia, [15] observed yield reduction due to contour bunds. The soil chemical and physical properties were not significantly influenced by practices. This is due to the fact that effects of these practices is long term [38] High maize and beans yields are due to the good fertility status of the soil. Most of the soil parameters were above the critical values, therefore adequate for plant growth. For banana, mulched plots had better yield reflecting the relative better nutrient conditions of the soils under mulch in terms of organic carbon, available phosphorus, extractable potassium and nitrogen.

5. CONCLUSIONS AND RECOMMENDATIONS

In light of the above results and discussion, it was concluded that soil loss was generally severe on maize and moderate on banana; while runoff was relatively low ranging between 1 and 3% on banana and maize intercropped with beans; respectively. The reduction in soil and runoff losses due to soil and water conservation practices was not significant during the period of the study. Nutrient losses were generally high on both systems and varied from one practice to another under maize but only K under banana. Nitrogen, phosphorus and potassium losses under banana varied from 46-107 Kg/ha/season; 23.6-48.1 Kg/ha/season; and 17 -53.9 Kg/ha/season; respectively. They ranged from 71-409 Kg/ha/season; 24-94 Kg/ha/season; 35-112 Kg/ha/season for N, P, and under maize; respectively. Based on the results and observations made in the study area there is need to improve the extension services in the micro-catchment, and the country at large, in order to promote and increase the adoption of the tested soil erosion control practices; and to conduct long term experiments to assess the efficiency of existing soil erosion control practices in the Lwiro micro-catchment and the region.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Brevik EC, Cerdà A, Mataix-Solera J, Pereg L, Quinton JN, Six J, Van Oost K. The interdisciplinary nature of soil. *SOIL*. 2015;1:117-129. DOI: 10.5194/soil-1-117-2015
2. Berendse F, van Ruijven J, Jongejans E, Keesstra S. Loss of plant species diversity reduces soil erosion resistance. *Ecosystems*. 2015;18(5):881-888. DOI: 10.1007/s10021-015-9869-6
3. Ligonja PJ, Shrestha RP. Soil erosion assessment in Kondoia eroded area in Tanzania using universal soil loss equation, geographic information systems and socioeconomic approach. *Land Degradation and Development*. 2015; 26(4):367-379. DOI: 10.1002/ldr.2215
4. Mekonnen M, Keesstra SD, Stroosnijder L, Baartman JEM, Maroulis J. Soil conservation through sediment trapping: A review. *Land Degradation and Development*. 2015;2015b:26(6):544-556. DOI: 10.1002/ldr.2308
5. Stanchi S, Falsone G, Bonifacio E. Soil aggregation, erodibility, and erosion rates in mountain soils (NW Alps, Italy). *Solid Earth*. 2015;6(2):403-414. DOI: 10.5194/se-6-403-2015
6. Majaliwa JGM, Twongyirwe R, Nyenje Oluca M, Ongom B, Sirike J, Mfitumukiza D, Azanga E, Natumanya R, Mwerera R, Barasa B. The effect of land cover change on soil properties around Kibale National Park in South Western Uganda; 2010.
7. Majaliwa JGM, Bashwira S, Tenywa M, Kansiime F. An overview of pollution loading into Lake Kivu basin. Third RUFORUM Biennial Meeting 24 - 28 September 2012, Entebbe, Uganda; 2012.
8. Giménez-Morera A, Ruiz Sinoga JD, Cerdà A. The impact of cotton geotextiles on soil and water losses from mediterranean rainfed agricultural land. *Land Degradation and Development*. 2010;21(2):210-217. DOI: 10.1002/ldr.971
9. Ligonja PJ, Shrestha RP. Soil erosion assessment in Kondoia eroded area in Tanzania using universal soil loss equation, geographic information systems and socioeconomic approach. *Land Degradation and Development*. 2015;26(4): 367-379. DOI: 10.1002/ldr.2215
10. Cerdà A, Jurgensen MF. The influence of ants on soil and water losses from an orange orchard in eastern Spain. *Journal*

- of Applied Entomology. 2007;132(4):306-314.
DOI: 10.1111/j.1439-0418.2008.01267.x
11. Seutloali KE, Beckedahl HR. Understanding the factors influencing rill erosion on road-cuts in the south eastern region of South Africa, *Solid Earth*. 2015;6: 633–641.
DOI: 10.5194/se-6-633-2015
 12. Ziegler AD, Negishi JN, Sidle RC, Gomi T, Noguchi S, Nik AR. Persistence of road runoff generation in a logged catchment in Peninsular Malaysia, *Earth Surf. Proc. Land*. 2015;32:1947–1970.
 13. Lasanta T, Cerdà A. Long-term erosional responses after fire in the Central Spanish Pyrenees: 2. Solute release. *Catena*. 2005; 60(1):81-100.
DOI: 10.1016/j.catena.2004.09.005
 14. Lieskovský J, Kenderessy P. Modelling the effect of vegetation cover and different tillage practices on soil erosion in vineyards: A case study in vrábce (Slovakia) using WATEM/SEDEM Land Degradation and Development. 2014;25(3):288-296.
DOI: 10.1002/ldr.2162
 15. Novara A, Gristina L, Saladino SS, Santoro A, Cerdà A. Soil erosion assessment on tillage and alternative soil managements in a Sicilian vineyard. *Soil and Tillage Research*. 2011;117:140-147.
 16. Mekonnen M, Keesstra SD, Baartman JE, Ritsema CJ, Melesse AM. Evaluating sediment storage dams: Structural off-site sediment trapping measures in northwest Ethiopia. *Cuadernos de Investigacion Geografica*. 2015a;41(1):7-22.
DOI: 10.18172/cig.2643
 17. Tenywa MM, Tukahirwa JB, Buruchara R, Adekunle AA, Mugabe J, Wanjiku C, Mutabazi S, Fungo B, Kashja NIM, Pali P, Mapatano S, Ngaboyisonga C, Farrow A, Njuki J, Abenakyo A. Agricultural innovation platform as a tool for development oriented research: Lessons and challenges in the formation and operationalisation. *Learning Publics Journal of Agriculture and Environmental Studies*. 2011;2(1):117-146.
 18. Ulimwengu J, Funes J, Headey D, You L. Paving the way for development? The impact of transport infrastructure on agricultural production and poverty reduction in the Democratic Republic of Congo. Washington DC: IFPRI Discussion Paper 00944, International Food Policy Research Institute (IFPRI). 2009;48.
 19. GTZ. Causes of soil degradation and development approaches to sustainable soil management; 1995.
 20. Zhang ZL. Human spoliation behaviors are the cause of northwest ecological degradation. *Renkou yanjiu [Demographic] [in Chinese.]* Costa, J.L., Aparicio, V., 1999;2:23–45.
 21. Prennushi G. Poverty reduction and the World Bank: Progress in Fiscal. 1999; 1998.
 22. Cohen AS, Bills R, Cocquyt CZ, Caljon AG. Impact of Sediment Pollution on Biodiversity in Lake Tanganyika. *Conservation Biology*. 1993;7(3):667-677.
 23. Donohue I, Irvine K, Wemaere A, Styles D, Hobbs W. Connecting land use with water quality: Sources, sinks and time bombs. Diffuse Pollution Conference, Dublin. *Water Resources Management*. 2003;56-60.
 24. Alin SR, O'Reilly CM, Dettman DL, Cohen AS, Palacios-Fest MR, McKee BA. Effects of land-use change on aquatic biodiversity: A view from the paleorecord at Lake Tanganyika, East Africa. *Geological Society of America*; 2002.
 25. Halbwachs M, Tietze K, Lorke A, Mudaheeranwa C. Investigations in Lake Kivu (East Central Africa) after the Nyiragongo eruption of January 2002. *Solidarités, Aide Humanitaire d'Urgence, Fparis*; 2002.
 26. Ischebeck O, Kabazimya K, Vilimumbalo S. Erosion a`Bukavu. *Proceedings Seminar 23– 28 January 1984, Bukavu, ISP Bukavu*. 1984;336.
 27. Lewis LA, Clay DC, DeJagher YMJ. Soil loss, agriculture and conservation in Rwanda: Toward sound strategies for soil management. *J. Soil Water Cons*. 1988; 418-421.
 28. Moeyersons J. La nature de l'érosion des versants au Rwanda. *Ann. Mus. roy. Afr. centr., Tervuren*. 1989;19.
 29. Moeyersons J, et al. A geomorphological assessment of landslide origin at Bukavu, Democratic Republic of the Congo. *Engineering Geology*. 2004;72:73-87.
 30. De Graaff J, Amsalu A, Bodnar F, Kessler A, Posthumus H, Tenge A. Factors influencing adoption and continued use of long-term soil and water conservation measures in five developing countries. *Applied Geogr*. 2008;28:271-280.
 31. Majaliwa et al. Efficiency of contour bunds in controlling soil, runoff and nutrient

- losses from major agricultural land-use types. In press; 2015.
32. Cahen L, Lepresonne J. The precambrian of the Congo, Rwanda and Burundi. In: Rankama K (ed). The Precambrian Interscience Publishers, New York. 1967; 143–290.
 33. Verhaeghe M. Inventaire des gisements calcaires, dolomies et travestins du Kivu, du Rwanda et du Burundi. Departement Geologie et Mines du Burundi. 1964;95.
 34. Ilunga L. Morphologie, volcanisme et sédimentation dans le rift du Sud-Kivu. Bulletin de la Société géographique de Liège. 1991;27:209-228.
 35. Pasteels P, Villeneuve M, De Paepe P, Klerkx J. Timing of the volcanism of the southern Kivu Province: Implications for the evolution of the western branch of the East African Rift system. Earth and Planetary Science Letters. 1989;94:353-363.
 36. Kampunzu AB, Vellutini PJ, Caron JP, Lubala RT, Kanika M,d Rumvegeri BT. Le volcanisme et l'évolution structurale du sud-Kivu. Un modele d'interpretation geodynamique du volcanisme distensif intracontinental. Bull. Cent. Rech. Explor. Prod. Eif_Aquitaine. 1983;7(1):257-271.
 37. Pouclet A. Le magmatisme du rift de l'Afrique Centrale, Interpretation des donnees chronologiques et relations structurales. 10eme colloque de Geologie Africaine, Resume, Montpellier. 1979;140-141.
 38. UNESCO. Lava series: Mapping the Hazardous Terrain using Remote Sensing, Geological Society Special Publication NO. 283 Majaliwa JGM, Tenywa MM, Rao KPC. 2015;2002.
 39. Soil Fertility in relation to landscape position and land use/covertypes: A case study of the lake kivu pilot learning site. Advances in Agriculture. 2015;Article ID 752936:8.
DOI: 10.1155/2015/752936
 40. Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis: A working manual; 2nd Edition. Sacred African Publishers, Nairobi, Kenya; 2002.
 41. Bray RH, Kurtz LT. Determination of total, organic, and available forms of phosphorus in soils. Soil Science. 1945;59:39-45.
 42. FAO. A provisional methodology for soil degradation assessment (with mapping of north Africa at a scale of 1: 5 million). FAO, Rome; 1979.
 43. FAO. Soil map of the World, Revised Legend. World Soil Resources Report.60. FAO, Rome; 1990.
 44. Majaliwa JGM. Effect of vegetation cover and biomass development on soil loss from maize based cropping systems. M.Sc thesis. Makerere University; 1998.
 45. Majaliwa JGM. Soil erosion from major agricultural land-use and their associated pollution loading in selected Lake Victoria micro-catchments. PhD thesis, Makerere University; 2004.
 46. Azanga E. Sediment and Nutrient Hotspot Areas Dynamics in Selected Micro-Catchment of the Lake Tanganyika Basin in Democratic Republic of Congo. Master Thesis, Makerere University, Kampala. 2013;97.
 47. Mulebeke R. Validation of a GIS-USLE model in a banana-based microcatchment of the Lake Victoria basin, Msc. Thesis, Makerere University; 2003.
 48. Stoorvogel JJ, Smaling EMA, Windmeijer PN. Calculating soil nutrient balances in Africa at different scales. I. Supranational scale. Fertility Research. 1993;35:227-237.
 49. Moreno-Ramón H, Quizembe SJ, Ibáñez-Asensio S. Coffee husk mulch on soil erosion and runoff: Experiences under rainfall simulation experiment. Solid Earth. 2014;5(2):851-862.
DOI: 10.5194/se-5-851-2014
 50. Cerdà A, Jurgensen MF. The influence of ants on soil and water losses from an orange orchard in eastern Spain. Journal of Applied Entomology. 2008;132(4):306-314.
DOI: 10.1111/j.1439-0418.2008.01267.x
 51. Tukahirwa J. Measurement, prediction and social ecology of soil erosion in Kabale, South western Uganda. PhD. Thesis, Institute of Environment and Resource Management, Makerere University; 1996.
 52. Landon JR. Tropical Soil Manual. Booker Agriculture. International Limited, Londres. 1995;450.
 53. Mulebeke R. Modeling soil loss in farming systems: Validation of a GIS based USLE in a banana-based microcatchment of Uganda's L. Victoria Basin, LAP Lambert Acad. Publ.; 2010.
 54. Le Bissonnais Y, Daroussin J, Jamagne M, Lambert JJ, Le Bas C, King D, Cerdan O, Le'onard J, Bresson LM, Jones RJA. Pan-

- European soil crusting and erodibility assessment from the European soil geographical database using pedotransfer rules. *Adv.*
55. Lal R. Soil erosion impact on agronomic productivity and environmental quality, *Crit. Rev. Plant Sci.* 1998;17:319–464.
 56. Simard RR, Beauchemin S, Haygarth P. M. Potential for preferential pathways of phosphorus transport. *J. Environ. Qual.* 2000;29:97–104.
 57. Ng Kee Kwong KF, Bholah A, Volcy L, Pynee K. Nitrogen and phosphorus transport by surface runoff from a silty clay loam soil under sugarcane in the humid tropical environment of Mauritius, *Agriculture, Ecosystems and Environment.* 2002;91:147–157.
 58. Fitzjohn C, Ternan JL, Williams AG. Soil moisture variability in a semi-arid gully catchment: Implications for runoff and erosion control. *Catena.* 1998;32:55–70.
 59. Meyles E, Williams A, Ternan L, Dowd J. Runoff generation in relation to soil moisture patterns in a small Dartmoor catchment, Southwest England, *Hydrol. Processes.* 2003;17:251–264.
 60. Castillo VM, Gomez-Plaza A, Martinez-Mena M. The role of antecedent soil water content in the runoff response of semi- arid catchments: A simulation approach, *J. Hydrol.* 2003;284:114–130.
 61. Wei LH, Zhang B, Wang MZ. Effects of antecedent soil moisture on runoff and soil erosion in alley cropping systems. *Agricultural Water Management.* 2007;94: 54–62.
 62. Martínez JRF, Zuazo VHD, Raya AM. Environmental impact from mountainous olive orchard under different soil-management systems (SE Spain). *Sci Total Environ.* 2006;358:46-60.
 63. Gómez JA, Guzmán MG, Giráldez JV, Fereres E. The influence of cover crops and tillage on water and sediment yield, and on nutrient, and organic matter losses in an olive orchard on a sandy loam soil. *Soil Till Res.* 2009;106:137-144.
 64. Farres PJ. The dynamics of rainsplash erosion and the role of soil aggregate stability. *Catena.* 1987;14:119–130.
 65. Lecomte V, Barriuso E, Bresson LM, Koch C, Le Bissonnais Y. Soil surface structure effect on isotoproturon and diflufenican loss in runoff. *J. Environ. Qual.* 2001;30:2113–2119.
 66. Lal M, Singh R. Carbon sequestration potential of Indian forests. *Environ Monit Assess.* 2000;60:315–327.
 67. Cerdà A. Soil physical quality changes under different management systems after 10 years in the Argentine humid Pampa. *Solid Earth.* 2015;6:361-371. DOI: 10.5194/se-6-361-2015

© 2018 Adidja et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/25118>*