

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/273995812>

# Genetic Relationships of Cassava Genotypes That are Susceptible or Tolerant to Cassava Brown Streak Disease in Uganda

Article in *Journal of Agricultural Science* · June 2013

DOI: 10.5539/jas.v5n7p107

CITATIONS

9

READS

418

10 authors, including:



Alex Abaca

MUNI University

7 PUBLICATIONS 106 CITATIONS

SEE PROFILE



R. S. Kawuki

National Agricultural Research Organization

68 PUBLICATIONS 1,656 CITATIONS

SEE PROFILE



Yona Baguma

NARO

33 PUBLICATIONS 909 CITATIONS

SEE PROFILE



Titus Alicai

National Agricultural Research Organization

208 PUBLICATIONS 2,623 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Disease diagnostics for sustainable cassava productivity in Africa [View project](#)



Improvement of nutritional value of African staple crops [View project](#)

## Genetic Relationships of Cassava Genotypes That are Susceptible or Tolerant to Cassava Brown Streak Disease in Uganda

Alex Abaca<sup>1,2</sup>, Robert Kawuki<sup>1</sup>, Phenihias Tukamuhabwa<sup>2</sup>, Yona Baguma<sup>1</sup>, Anthony Pariyo<sup>1</sup>, Titus Alicai<sup>1</sup>,  
C. Christopher Omongo.A<sup>1</sup>, Phillips Abidrabo<sup>1</sup>, Kasifa Katono<sup>1</sup> & Anton Bua<sup>1</sup>

<sup>1</sup>National Crops Resources Research Institute, Namulonge, Kampala, Uganda

<sup>2</sup>Makerere University Kampala, Kampala, Uganda

Correspondence: Alex Abaca, National Crops Resources Research Institute, Namulonge, P.O. Box 7084, Kampala, Uganda. Tel: 256-0-782-285-955. E-mail: abal-2007@hotmail.com

Received: June 4, 2011 Accepted: September 13, 2012 Online Published: June 15, 2013

doi:10.5539/jas.v5n7p107

URL: <http://dx.doi.org/10.5539/jas.v5n7p107>

### Abstract

A total of 99 cassava genotypes whose field reaction to cassava brown streak disease (CBSD) was known, were assayed with 30 simple sequence repeat (SSR) markers to establish their genetic parallels. Two categories of CBSD reaction were considered: CBSD-susceptible genotypes (characterized by > 60% root and foliar CBSD incidence), and CBSD-tolerant genotypes (characterized by < 15% root and foliar CBSD incidence). DNA was extracted from leaf samples using the miniprep method and genotyped using ABI 3730 DNA sequencer. The test genotypes at 0.02 similarity coefficient, CBSD-tolerant and CBSD-susceptible genotypes clustered into 5 main sub-clusters. When data were subjected to principle component analysis (PCA) irrespective of the CBSD reaction grade, the first three principal components accounted for 68% of the total genetic variation. Despite having different number of individuals, the observed heterozygosity ( $H_o$ ) for CBSD-susceptible ( $H_o = 0.58$ ) and CBSD-tolerant ( $H_o = 0.66$ ) were comparable and provide scope for long-term CBSD breeding and/or gene tapping.

**Keywords:** Cassava brown streak virus, cluster analysis, foliar symptoms, simple sequence repeat marker

### 1. Introduction

Genetic diversity is essential for genetic advancement of crops and animals. For crops, by exploring the level and distribution of genetic diversity as well as the evolutionary processes shaping this diversity, plant breeders can come out with new improved varieties. Moreover, the analysis of genetic variation helps in the choice of germplasm for breeding purposes. Correct genetic analysis can lead to the identification of new combinations with maximum genetic variability, which can then be used for introgression of desirable genes into the available genetic base (Smith, 1984; Cox, Murphy, & Rodgers, 1986; Mohammadi & Prasanna, 2003).

Variation in allele frequency at many unlinked loci is the preferred way to assess genetic diversity and/or genetic differentiation (Nei, 1973; Wu & Tansley, 1993). Indeed, molecular markers have been used to quantify genetic diversity in a number of crops including: sorghum (Dj'e, Heuertz, lef'ebvre, & Vekemans, 2000); beans (Cabral et al. (2011); Foster et al. (2010); rice (Vanniarajan, Vinod, & Pereira, 2012); cotton (Nagaraj Basavaraddi, 2007); and finger millet (Kumari & Pande, 2010) amongst others. For cassava genetic diversity studies have been conducted both in Latin America (Elias, Paraud & Robert, 2000; Elias, Penet, Vindry, MacKey, Panaud & Robert, 2001; Elias, Mühlen, Mckey, Roa, & Tohme, 2004) and in Africa (Mkumbira et al., 2003; Fregene et al., 2003). It suffices to note that genetic diversity of cassava has been evaluated using different molecular markers: restriction fragment length polymorphisms (RFLPs) (Beeching et al., 1993); random amplified polymorphic DNA (RAPD) markers (Marmey, Beeching, Hamon, & Charrier, 1994); simple sequence repeat markers (SSR) markers (Fregene et al., 2003), and single nucleotide polymorphism (SNPs) (Lopez et al., 2005; Kawuki, Ferguson, Labuschagne, Herselman, & Kim, 2009). In cassava, molecular markers have been used to study: (i) relationship between cassava and its wild relatives (Roa et al., 1997); (ii) intra-specific variation to enable the identification of duplicates (Beeching et al., 1993); (iii) understand genetic structure in reference to ecological adaptation (Bornierbale, Maya, Claros, & Iglesias, 1995), and (iv) resistance and/or susceptibility to diseases (Sanchez et al., 1999; Fregene et al., 2003). In this paper we are using SSR markers to help discern the genetic relationships

between CBSD-susceptible and CBSD-tolerant genotypes in order to efficiently design a durable CBSD breeding programme.

We have selected CBSD because it's one of the 100 most dangerous diseases in the world (Donald, 2010). CBSD is caused by *cassava brown streak virus* (CBSV) (Monger, Seal, Issac, & Foster, 2001) and *Uganda cassava brown streak virus* (UCBSV), both of which, are prevalent in Uganda (Alicai Titus Personal Information). For over 70 years CBSD had been known to be confined to low altitude areas, especially in the coastal areas of Kenya, Tanzania and Mozambique. The presence of CBSD at high altitude areas including Uganda is just recent (Alicai et al., 2007). The disease causes characteristic above and below ground symptoms (Nichols, 1950; Hillocks & Thresh, 2000; Hillocks & Jennings, 2003), and as of this writing no cassava variety is resistant to it. Only a handful of tolerant varieties have been identified through field evaluations in hotspot areas (Abaca et al., 2012).

Technically, utilization of different and diverse CBSD tolerant lines would be desirable for durable CBSD breeding. Whether or not the presumed CBSD tolerate lines have a shared pedigree is unknown. It is also interesting to know the genetic relationships between the CBSD-susceptible and/or CBSD-tolerant genotypes. Thus, the overall objectives of this study were to determine the genetic relationships between CBSD-susceptible and/or CBSD-tolerant cassava genotypes using the simple sequence repeats markers.

## 2. Materials and Methods

### 2.1 Plant Material and DNA Extraction

Fresh leaf samples were collected from 100 cassava genotypes whose field reaction to CBSD was known; these materials had previously been evaluated for reaction to CBSD at Namulonge, which is a hotspot for both strains of brown streak strains (Abaca et al., 2012). However, one genotype had a lot of missing values and was omitted from the SSR analysis. The CBSD field reaction of these genotypes is presented in Table 1. DNA was extracted using the miniprep protocol as described by (Ranjana, Morag, Melaku, Dominique and Ivan (2009).

Table 1. Field reaction of the 100 test cassava genotypes to cassava brown streak disease in Uganda

Genotype	Reaction to CBSD	Genotype	Reaction to CBSD	Genotype	Reaction to CBSD	Genotype	Reaction to CBSD
<b>Elite</b>		B134	S	Kyembabazi	S	Nasiwa	S
NASE 11	S	TME 05	S	Njule silverbrown	S	Mweru	S
MH95/0402	S	MM02/0258	S	Kindimupale	S	Ntyaraboke tall	S
95/SE-00094	S	MM01/1003	S	Nyapatek	S	Gwalanda	S
MM01/1457	S	NASE 12	S	Kalangwa	S	Mpologoma	S
Silira 15 B2	S	MH04/300	S	Kyemigisha	S	Nyapamitu	S
95/SE-0088	S	MH04/0486	S	Bukalasa	S	Bukalasa 8	S
TZ06/130	T	MH04/0479	S	Siira	S	Rutangi	S
95/SE-00036	S	MM06/400	S	Masindi 2	S	Esau	S
TZ06/140	S	TME 204	S	Kakofira	S	Kalintunsi	S
MM06/0532	S	/0427	S	Kalisa	S	Birungi	S
MM06/529	S	349 KAK	S	Tamale	S	Ssesubiri	S
I92/00067	S	MM96/0686	T	Kabiriti	S	Serubola	S
MM97/2358	S	NASE 1	T	Abigaba	S	Rusidi	S
NASE 4	S	CR 20A - 1	T	Kirimupale	S	Kasule	S
MH04/1636	S	MH04/0042A	S	Alado alado	S	Kabwa	S
TME 14	S	266 BAM	S	Tim Tim	S	Luhujo white	S
I92/00057	S	MH97/2961	S	Ntangali	S	Bamunanika	S
96/1630	S	<b>Local</b>		Matooke	S	Kibao	S
MH04/0016	S	Nyarunega	S	Mukuma	S	Lukiru	S
MH04/2018	S	MM96/0876	S	Masindi 3	S	Kamuhanda	S
MM02/0169	S	Njule white	S	Kabagambe	S	Kiberu	S
MM02/1970	S	Njule zigzag	S	Kisembo	S	Bao	S
NASE 3	S	Nyamatia	S	Masindi 1	S	Rwamutere	S
MM98/3055	S	Kamia	S	Sentongo	S		
MM96/4271	T	Njule red	S	Lugujo brown	S		

Susceptible genotypes characterized by > 60% root and foliar CBSD incidence; tolerant genotypes characterized by < 15% root and foliar CBSD incidence.

## 2.2 DNA Genotyping

Polymerase chain reaction (PCR) amplifications with 30 SSR primers (Table 2) were performed in 10  $\mu$ l reactions that contained 10 ng of DNA, 1 mole of each primer, 1x *Taq* polymerase buffer, 2 mM  $MgCl_2$ , 0.2 mM deoxynucleotide triphosphates (dNTPs) and 0.375 U *Taq* polymerase (New England Biolabs Inc.). The PCR profile was 95°C for 2 min, followed by 30 cycles of 95°C for 30 sec, 45-55°C for 1 min and 72°C for 1 min and a final extension at 72°C for 30 min. Based on previous studies, these 30 SSR markers were chosen because of their diallelic nature, reproducibility and distribution across the genome (Moyib, Odunola, & Dixon, 2007; Raghu et al., 2007). Aliquots of 1  $\mu$ l of the PCR products were added to 9  $\mu$ l of the master-mix that contained HiDi formamide and GeneScan-500 LIZ size standard (1 ml of HiDi + 12  $\mu$ l of 500-LIZ). Amplicons were denatured (95°C for 3 min) and subjected to capillary electrophoresis using the ABI 3730 DNA sequencer (Applied Biosystems) and allele calls made using the GeneMapper ver. 3.7 software (Applied Biosystems). All genotyping work was done at the Biosciences eastern and central Africa (BeCA) hub in Nairobi, Kenya.

Table 2. SSR locus, genetic map location, primer sequences, and annealing temperature of markers used for genotyping

SSR locus	Genetic Map Location	Left primer (Reverse)	Right primer (Forward)	Product Size (bp)	T (°C)
SSRY9	D	ACAATTCATCATGAGTCATCAACT	CCGTTATTGTTCTGCTCCT	278	55
SSRY12	I	AACTGTCAAACCATTCTACTTGC	GCCAGCAAGGTTTGCTACAT	266	55
SSRY19	K	TGTAAGGCATTCCAAGAATTATCA	TCTCCTGTGAAAAGTGCATGA	214	55
SSRY20	R	CATTGGACTTCTACAAATATGAAT	TGATGGAAAAGTGTTATGTCCTT	143	55
SSRY21	N	CCTGCCACAATATTGAAATGG	CAACAATTGGACTAAGCAGCA	192	55
SSRY34	F	TTCCAGACCTGTTCCACCAT	ATTGCAGGGATTATTGCTCG	279	55
SSRY38	G	GGCTGTTCTGTGATCCTTATTAAC	GTAGTTGAGAAAACCTTTCATGAG	122	55
SSRY51	E	AGGTTGGATGCTTGAAGGAA	GGATGCAGGAGTGCTCAACT	298	55
SSRY52	I	GCCAGCAAGGTTTGCTACAT	AACTGTCAAACCATTCTACTTGC	266	55
SSRY59	M	GCAATGCAGTGAACCATCTTT	CGTTTGTCTTTCTGATGTTT	158	55
SSRY63	nd	TCAGAATCATCTACCTTGGCA	AAGACAATCATTTTGTGCTCCA	290	55
SSRY64	H	CGACAAGTCGTATATGTAGTATTCACG	GCAGAGGTGGCTAACGAGAC	194	55
SSRY69	E	CGATCTCAGTCGATACCCAAG	CCTCCGTTGCAGGCATTA	239	55
SSRY82	O	TGTGACAATTTTCAGATAGCTTCA	CACCATCGGCATTAACCTTTG	211	55
SSRY100	P	ATCCTTGCTGACATTTTGC	TTCGAGAGTCCAATTGTTG	210	55
SSRY102	F	TTGGCTGCTTCTACTAATGC	TTGAACACGTTGAACAACCA	179	55
SSRY103	B	TGAGAAGGAAACTGCTTGAC	CAGCAAGACCATCACCAGTTT	272	55
SSRY105	G	CAAACATCTGCACTTTTGGC	TCGAGTGGCTTCTGGTCTTC	225	55
SSRY106	B	GGAAACTGCTTGCAAAAGA	CAGCAAGACCATCACCAGTTT	270	55
SSRY108	D	ACGCTATGATGTCCAAAGGC	CATGCCACATAGTTCGTGCT	203	55
SSRY110	X	TTGAGTGGTGAATGCGAAAG	AGTGCCACCTTGAAAGAGCA	247	55
SSRY135	G	CCAGAAACTGAAATGCATCG	AACATGTGCGACAGTGATTG	253	45
SSRY161	E	AAGGAACACCTCTCCTAGAATCA	CCAGCTGTATGTTGAGTGAGC	220	55
SSRY164	H	TCAAACAAGAATTAGCAGAACTGG	TGAGATTCGTAATATTCATTCACTT	187	45
SSRY171	C	ACTGTGCCAAAATAGCCAAATAGT	TCATGAGTGTGGGATGTTTTATG	291	55
SSRY177	U	ACCACAAACATAGGCACGAG	CACCCAATTCACCAATTACCA	268	45
SSRY179	nd	CAGGCTCAGGTGAAGTAAAGG	GCGAAAGTAAGTCTACAACCTTTCTAA	226	55
SSRY180	E	CCTTGGCAGAGATGAATTAGAG	GGGGCATTCTACATGATCAATAA	163	55
SSRY181	K	GGTAGATCTGGATCGAGGAGG	CAATCGAAACCGACGATACA	199	55
SSRY182	Q	GGAATTCTTGCTTATGATGCC	TTCTTTACAATTCTGGACGC	253	55

nd= Genetic location not known, and T= temperature in degree Celsius.

## 2.3 Data Analysis

The generated diploid genetic data was used to generate allelic frequency and dissimilarity matrices using GeneAlex software, ver. 6.4 (Peakall & Smouse, 2006). Further, cluster analyses were performed to measure the degree of relatedness between accessions in the DARwin5 software, ver. 5.0.158 (Perrier & Jacquemond, 2006) based on the UPGMA method.

### 3. Results

#### 3.1 Genetic Relationships between CBDS-Tolerant and CBDS-Susceptible Cassava Genotypes

A total of 30 SSR loci were analyzed in 99 cassava genotypes classified as either CBDS-susceptible (94 genotypes) or CBDS-tolerant (five genotypes). As of this writing we only have five cassava genotypes that can be classified as tolerant i.e., with CBDS root and foliar incidence  $\leq 15\%$ . Most of the cassava varieties available in the Ugandan germplasm are highly susceptible to CBDS. The dendrogram of genetic similarity of the test genotypes with the coefficient of genetic similarity ranging from 0 to 0.1, is presented in Figure 1. Two main grouping were apparent, with only one CBDS-tolerant genotype (NASE 1) being assigned to one cluster and the other four CBDS-tolerant genotypes (CR 20A -1, MM96/4271, MM96/0686 and TZ 06/130) being assigned to the second cluster. However, at 0.02 similarity coefficient, the genotypes clustered into 5 main sub-clusters. The first sub-cluster consisted of 40 genotypes including the four CBDS-tolerant genotypes (Figure 1). The second sub-cluster consisted of only five CBDS-susceptible genotypes. The third sub-cluster consisted of 51 genotypes of which only NASE 1 was CBDS-tolerant. The fourth and fifth sub-clusters consisted respectively of two and one CBDS- susceptible genotypes.




Figure 1. UPGMA dendrogram of 99 cassava genotypes based on the dissimilarity matrix calculated from the SSR markers. The arrow show the position of the CBDS-tolerant genotypes (NASE 1, CR20A-1, MM96/0686, MM96/4271 and TZ06/130)




Figure 2. Principle component analysis of 99 cassava genotypes analyzed using 30 SSR markers

### 3.2 Genetic Diversity of Elite and Local Cassava Genotypes

When the genotypic data were re-analyzed with cassava classified as either local (cassava commonly grown by farmers and not resulting from a formal breeding process) or elite (cassava resulting from a formal breeding process), the results showed that local genotypes had relatively more allelic diversity than their elite counterparts (Table 3). Observed heterozygosity ( $H_o$ ) was also high in all the two classification with the averages of  $0.61 \pm 0.04$  and  $0.62 \pm 0.004$  for the local and elite genotypes respectively (Table 3). Similarly, expected heterozygosity ( $H_e$ ) was high within the local genotypes than the elite (Table 3). Analysis of molecular variance (AMOVA) revealed that 98% of the genetic variation was distributed among individual genotypes and not groups that is, between elite and local genotypes. Results from principle component analysis showed similar groupings with the majority of the elite genotypes somewhat clustering separately from the local genotypes (Figure 2). The first three principal components accounted for 38.02%, 15.59% and 14.84 % of the total genetic variation, respectively.

Table 3. Number of alleles per SSR locus and genetic diversity parameters for local and elite cassava genotypes

Populations	Sample size (n)	Na	Ne	$H_o$	$H_e$	UHe
Local	54	$5.1 \pm 0.32$	3.13	$0.61 \pm 0.04$	$0.61 \pm 0.03$	0.61
Elite	45	$4.6 \pm 0.29$	2.89	$0.62 \pm 0.04$	$0.58 \pm 0.03$	0.59
<b>Mean</b>	<b>49.5</b>	<b><math>4.9 \pm 0.31</math></b>	<b>3.01</b>	<b><math>0.62 \pm 0.04</math></b>	<b><math>0.6 \pm 0.03</math></b>	<b>0.601</b>

Na=Average number of different alleles, Ne=Average number of effective alleles,  $H_o$ = average observed heterozygosity,  $H_e$ =Average expected heterozygosity and UHe= Average unbiased expected heterozygosity.

#### 4. Discussion

The major objective of this study was to establish whether or not the CBSD-Tolerant and CBSD-susceptible genotypes are genetically similar. The CBSD-tolerant genotypes (NASE 1, CR 20A -1, MM96/4271, MM96/0686 and TZ 06/130) were placed in two separate clusters; one cluster had only one genotype (NASE 1) and the other cluster had the remaining four genotypes. NASE 1 also referred to as TMS 60142 is one of the Tropical Manihot Selections that were derived from clone 58308 (a third backcross) at Amani, Tanzania, where the pioneer CBSD breeding efforts were undertaken. This genotype was officially released in Uganda in the early 1990s owing to its resistance to CMD, the key viral constraint then. On the other hand, both MM96/4271 and MM96/0686 are genotypes from the International Institute of Tropical Agriculture (IITA) breeding programme; TZ06/130 was selected from open pollinated seeds sourced from CBSD-tolerant genotypes in Tanzania; and CR20A-1 is a Latin American genotype sourced from International Centre for Tropical Agriculture (CIAT). Basing on these results, it's likely that we are dealing with different sources of CBSD tolerance, which appears to be consistent with the aforementioned historical pedigree information. It is therefore envisaged that combining genotypes from the different clusters and/or sub-clusters as parents in the breeding programme would result into diversifying and/or increasing the dosage effects required for durable CBSD resistance breeding.

The implications of this study are of direct relevance towards the management and control of the spread of CBSD, and in breeding for resistance. Several surveys conducted in East and Central Africa (Legg & Pheneas, 2007) have shown that there are severe forms of the CBSD due to cassava brown streak virus (CBSV) and Uganda cassava brown streak virus (UCBSV), and that the disease has spread to neighbouring countries such as Democratic Republic of Congo, Rwanda, Burundi amongst others. In managing the spread of the disease, cultivating tolerant genotypes has been demonstrated as the most effective means (ASERECA, 2009). In the absence of resistant varieties, it's these tolerant varieties that can be multiplied and thereafter distributed to farmers. However, because they are not immune to CBSVs, their usefulness will be short-lived as a result of accumulation of virus with each cycle of propagation. Nonetheless, these tolerant genotypes should be able to provide descent yields even with the fifth cycle of propagation after which, virus elimination should be done using tissue culture techniques.

In parallel, the identified tolerant genotypes can be utilized to initiate a breeding scheme that pyramids CBSD tolerance alleles. Inter-mating CBSD tolerant cassava genotypes which have other desirable agronomic and/or consumer quality traits, from the different sub-clusters would ensure the diversification of tolerance to the disease while creating new genotypes. Preliminary information on the genetics of inheritance for CBSD indicate that both additive and non-additive gene effects are important (Munga, 2008) and thus justifying the need for recurrent selection for CBSD breeding. With this in mind, the tolerate lines can be crossed in different combinations to generate progeny for further evaluation and hybridization. In addition, by combining different genes that relate to different sources of resistance, epistatic interaction may be identified such that higher levels of resistance can be developed.

High levels of heterozygosity were observed in the cassava genotypes, with an average of 0.611 for both local and elite cassava genotypes. These high levels of heterozygosity can be attributed to the out-crossing nature of cassava. This is similar to the explanation of Karuri, Ateka, Amata, Nyende and Muigai (1998) that self incompatibility in the flowers of sweet potatoes results in allogamy, thus, increasing genetic heterozygosity. Furthermore, despite some differences in allelic diversity and the observed clustering, most variation was detected within individuals, which accounted for 98% of the total genetic variation, a finding which further confirms the heterozygous nature of cassava.

This is the first methodical analysis of genetic relationships between CBSD-susceptible and CBSD-tolerant cassava genotypes in Uganda. Our findings indicate that the current sources of CBSD tolerance are distinct, which is desirable when dealing with viral pathogens that are highly variable and/or have potential to mutate. The tolerant genotypes can either be hybridized amongst themselves to concentrate favorable alleles and/or selfed to fix the desirable alleles.

#### Acknowledgement

The authors thank the Millennium Science Initiative (MSI) that provided financial support through the Uganda National Council for Science and technology (UNCST). We are grateful to the technicians of the National Cassava programme for their support towards data collection and trial management.

## References

- Abaca, A., Kawuki, R., Tukamuhabwa, P., Baguma, Y., Pariyo, A., Alicai, T., ... Bua, A. (2012). Evaluation of Local and Elite Cassava Genotypes for Resistance to Cassava Brown Streak Disease (CBSD) in Uganda. *Journal of Agronomy*, 11(3), 65-72. <http://dx.doi.org/10.3923/ja.2012.65.72>
- Alicai, T., Omongo, C. A., Maruthi, M. N., Hillocks, R. J., Baguma, Y., Kawuki, R., ... Colvin, J. (2007). Re-emergence of cassava brown streak disease in Uganda. *Plant Disease*, 91, 24-29. <http://dx.doi.org/10.1094/PD-91-0024>
- ASERECA. (2009). *Communication and training resource kit for the awareness campaign on cassava brown streak disease*. Staple Crops Programme, Plot 5 Mpigi Road, P.O Box 765, Entebbe – Uganda.
- Beeching, J. R., Marmey, P, Gavaldà, M. C., Noirot, M., Hayson, H. R., Hughes, M. A., & Charrier, A. (1993). An assessment of genetic diversity within a collection of cassava (*Manihot esculenta*, Crantz) germplasm using molecular markers. *Annals of Botany*, 72(6), 515-571. <http://dx.doi.org/10.1006/anbo.1993.1139>
- Bonierbale, M. W., Maya, M. M., Claros, J. L., & Iglesias, C. (1995). Application of molecular markers to describing the genetic structure of cassava gene pools. In *Proceedings of the Second International Scientific Meeting of the Cassava Biotechnology Network, Bogor, Indonesia, 22-26 August 1994, Volume I. CIAT Working document No. 150. CIAT, Cali, Colombia* (pp 106-112).
- Cabral, P. D. S., Soares, T. C. B., Lima, A. B. P., de Miranda, F. D., Souza, F. B., & Gonçalves, L. S. A. (2011). Genetic diversity in local and commercial dry bean (*Phaseolus vulgaris*) accessions based on microsatellite markers. *Genetics and Molecular Research*, 10(1), 140-149. <http://dx.doi.org/10.4238/vol10-1gmr993>
- Cox, T., Murphy, J., & Rodgers, D. (1986). Changes in genetic diversity in the red winter wheat regions of the United States. *Proceedings of the National Academy of Science (USA)*, 83, 5583-5586. <http://dx.doi.org/10.1073/pnas.83.15.5583>
- Dj`e, Y., Heuertz, M., Lef`ebyre, C., & Vekemans, X. (2000). Assessment of genetic diversity within and among germplasm accessions in cultivated sorghum using microsatellite markers. *Theoretical and Applied Genetics*, 100, 918–925. <http://dx.doi.org/10.1007/s001220051371>
- Donald, G. M. J. (2010). *Virus Ravages Cassava Plants in Africa*. The New York Times, Published on May 31, 2010. Retrieved 11th October, 2011, from <http://www.nytimes.com/2010/06/01/science/01cassava.html>
- Elias, M., Mühlen, G. S, McKey, D., Roa, A. C., Tohme, J. (2004). Genetic diversity of traditional South American landraces of cassava (*Manihot esculenta* Crantz): an analysis using microsatellites. *Economic Botany*, 58, 242-256. [http://dx.doi.org/10.1663/0013-0001\(2004\)058\[0242:GDOTSA\]2.0.CO;2](http://dx.doi.org/10.1663/0013-0001(2004)058[0242:GDOTSA]2.0.CO;2)
- Elias, M., Paraud, O., Robert, T. (2000). Assessment of genetic variability in a traditional cassava (*Manihot esculenta* Crantz) farming system using AFLP markers. *Heredity*, 85(3), 219-230. <http://dx.doi.org/10.1046/j.1365-2540.2000.00749.x>
- Elias, M., Penet, L., Vindry, P., MacKey, D., Panaud, O., & Robert, T. (2001). Unmanaged sexual reproduction and the dynamics of genetic diversity of a vegetatively propagated crop plant, cassava (*Manihot esculenta* Crantz), in a traditional farming system. *Molecular Ecology*, 10, 895-907. <http://dx.doi.org/10.1046/j.0962-1083.2001.01331.x>
- Schork, N. J., Fallin, D., & Lanchbury, J. S. (2000). Single nucleotide polymorphisms and the future of genetic epidemiology. *Clinical genetics*, 58(4), 250-264. <http://dx.doi.org/10.1186/1471-2229-10-13>
- Fregene, M., Suarez, M., Mkumbira, J., Kulembeka, H., Ndedya, E., Kulaya, A., ... Kresovich, S. (2003). Simple sequence repeat marker diversity in cassava landraces: genetic diversity and differentiation in an asexually propagated crop. *Theoretical and Applied Genetics*, 107, 1083-1093. <http://dx.doi.org/10.1007/s00122-003-1348-3>
- Hillocks, R. J., & Thresh, M. J. (2000). Cassava mosaic and cassava brown streak virus- diseases in Africa. *Roots*, 7, 1-12. <http://dx.doi.org/10.1080/0967087031000101061>
- Hillocks, R. J., & Jennings, D. L. (2003). Cassava brown streak disease: a review of present knowledge and research needs. *International Journal of Pest Management*, 49, 225-234.
- Karuri, H. W., Ateka, E. M., Amata, R., Nyende, A. B., & Muigai, A. W. T. (1998). Characterization of Kenyan sweet potato genotypes for SPVD resistance and high dry matter content using simple sequence repeat markers. *African Journal of Biotechnology*, 8(10), 2169-2175.



- Kawuki, R. S., Ferguson, M., Labuschagne, M., Herselman, L., & Kim, D. (2009). Identification, characterisation and application of single nucleotide polymorphisms for diversity assessment in cassava (*Manihot esculenta* Crantz). *Molecular Breeding*, *23*, 669-684. <http://dx.doi.org/10.1007/s11032-009-9264-0>
- Kumari, K., & Pande, A. (2010). Study of genetic diversity in finger millet (*Eleusine coracana* L. Gaertn) using RAPD markers. *African Journal of Biotechnology*, *9*(29), 4542-4549. Retrieved from <http://www.academicjournals.org/AJB>
- Legg, J. P., & Pheneas, N. (2007). New spread of cassava brown streak virus disease and its implications for the movement of Germplasm in the East and Central African Region. *International Journal of Pest Management*, *53*, 179-182.
- Lopez, C., Piegu, B., Cooke, R., Delsen, M., Tohme, J., & Verdier, V. (2005). Using cDNA and genomic sequences as tools to develop SNP strategies in cassava (*Manihot esculenta* Crantz). *Theoretical and Applied Genetic*, *110*, 425-431. <http://dx.doi.org/10.1007/s00122-004-1833-3>
- Marmey, P., Beeching, J. R., Hamon, S., & Charrier, A. (1994). Evaluation of cassava (*Manihot esculenta* Crantz) germplasm collection using RAPD markers. *Euphytica*, *14*, 203-209. <http://dx.doi.org/10.1007/BF00040402>
- Mkumbira, J., Chiwona-Karlton, L., Lagercrantz, U., Mahungu, N. M., Saka, J., Mhone, A., ... Rosling, H. (2003). Classification of cassava into 'bitter' and 'cool' in Malawi: from farmers' perception to characterisation by molecular markers. *Euphytica*, *132*, 7-22. <http://dx.doi.org/10.1023/A:1024619327587>
- Mohammadi, S., & Prasanna, B. (2003). Review & interpretation. Analysis of genetic diversity in crop plants. *Crop Science*, *43*, 1235-1248. <http://dx.doi.org/10.2135/cropsci2003.1235>
- Monger, W. A., Seal, S., Issac A. M., & Foster, G. D., (2001a). Molecular characterization of the cassava brown streak virus coat protein. *Plant pathology*, *50*, 527-534. <http://dx.doi.org/10.1046/j.1365-3059.2001.00589.x>
- Moyib O. K., Odunola O. A., Dixon A. G. O. (2007). SSR markers reveal genetic variation between improved cassava cultivars and landraces within a collection of Nigerian cassava germplasm. *African Journal of Biotechnology*, *6*(23), 2666-2674.
- Munga Theresia Luvuno. (2008). *Breeding for Cassava Brown Streak Resistance in Coastal Kenya*. PhD thesis, Faculty of Science and Agriculture, University of KwaZulu-Natal, Republic of South Africa.
- Nagaraj, B. (2007). *Genetic diversity studies in Gossypium hirsutum L. Cotton, for Yield contributing characters, Fibre traits and in Vitro Regeneration*. MSc Thesis. University of Agricultural Sciences, Dharwad.
- Nei, M. (1973). Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Science*, *70*, 3321-3323. <http://dx.doi.org/10.1073/pnas.70.12.3321>
- Nichols, R. F. W. (1950). The brown streak disease of cassava: distribution climatic effects and diagnostic symptom. *East African Agriculture*, *15*, 154-160.
- Peakall, R., & Smouse, P. E. (2006). GENALEX 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, *6*, 288-295. <http://dx.doi.org/10.1111/j.1471-8286.2005.01155.x>
- Perrier, X., & Jacquemoud-Collet, J. P. (2006). DARwin Software. Retrieved May 15th, 2011, from <http://darwin.cirad.fr/darwin>
- Raghu, D., Senthil, N., Saraswathi, T., Raveendran, M., Gnanam, R., Venkatachalam, R., ... Mohan, C. (2007). Morphological and simple sequence repeats (SSR) based fingerprinting of South Indian cassava germplasm. *International Journal of Integrative Biology*, *1*(2), 141-148.
- Ranjana, B., Morag, F., Melaku, G., Dominique, D., & Ivan, I. (2009). Field collection, preservation and large scale DNA extraction procedures for cassava (*Manihot esculenta* Crantz). *African Journal of Biotechnology*, *8*(15), 3424-3430.
- Roa, A., Maya, M., Duque, M., Tohme, J., Allem, A., & Bonierbale, M. (1997). AFLP analysis of the relationships among cassava and other *Manihot* species. *Theoretical and Applied Genetics*, *95*, 741-750. <http://dx.doi.org/10.1007/s001220050620>
- Sanchez, G., Restrepo, S., Duque, M., Fregene, M., Bonierbale, M. W., & Verdier, V. (1999). AFLP assessment of genetic variability in cassava accessions (*Manihot esculenta* Crantz) resistant and susceptible to the cassava bacterial blight (CBB). *Genome*, *42*, 163-172. <http://dx.doi.org/10.1139/g98-124>

- Smith, J. (1984). Genetic variability within U.S. hybrid maize: multivariate analysis of isozyme data. *Crop Science*, 24, 1041-1046. <http://dx.doi.org/10.2135/cropsci1984.0011183X002400060009x>
- Vanniarajan, C., Vinod, K. K., & Pereira, A., (2012). Molecular evaluation of genetic diversity and association studies in rice (*Oryza sativa* L.). *Journal of Genetics*, 91, 9-19. <http://dx.doi.org/10.1007/s12041-012-0146-6>
- Wu, K. S., & Tanksley, S. D. (1993). Abundance, polymorphism and genetic mapping of microsatellites in rice. *Molecular Genetics and Genomics*, 241, 225-235.

### Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).